



Diagnosing pancreatic cancer in general practice: a cross-sectional study on associations between suspicion of cancer, urgent referral and time to diagnosis

Line F. Virgilsen, Linda A. Rasmussen, Peter Vedsted & Henry Jensen

To cite this article: Line F. Virgilsen, Linda A. Rasmussen, Peter Vedsted & Henry Jensen (2022) Diagnosing pancreatic cancer in general practice: a cross-sectional study on associations between suspicion of cancer, urgent referral and time to diagnosis, Scandinavian Journal of Primary Health Care, 40:1, 78-86, DOI: [10.1080/02813432.2022.2036491](https://doi.org/10.1080/02813432.2022.2036491)

To link to this article: <https://doi.org/10.1080/02813432.2022.2036491>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 11 Feb 2022.



[Submit your article to this journal](#)



Article views: 1361



[View related articles](#)



[View Crossmark data](#)



Citing articles: 2 [View citing articles](#)

Diagnosing pancreatic cancer in general practice: a cross-sectional study on associations between suspicion of cancer, urgent referral and time to diagnosis

Line F. Virgilsen^a, Linda A. Rasmussen^a, Peter Vedsted^{a,b} and Henry Jensen^a

^aResearch Centre for Cancer Diagnosis in Primary Care, Research Unit for General Practice, Aarhus, Denmark; ^bDepartment of Clinical Medicine, Aarhus University, Aarhus C, Denmark

ABSTRACT

Objective: This study aimed to investigate the first point of contact in patients diagnosed with pancreatic cancer, and to study factors associated with the GP's suspicion of cancer, Cancer Patient Pathway (CPP) referral and long diagnostic interval.

Design: Cross-sectional study combining register and survey data.

Patients: Patients with incident pancreatic cancer recorded in the Danish National Patient Register (n = 303).

Main outcome measures: The patient's first point of symptoms presentation, GP's cancer suspicion, CPP referral and diagnostic interval.

Results: General practice was the first point of contact for 85.5% of the population. At the first consultation, cancer was suspected in 32.7% and 22.9% were referred to a CPP. The GPs were more likely to suspect cancer or serious illness in patients aged >70 years (prevalence rate ratio (PRR) 1.34, 95% CI 1.09–1.66) and among patients with high comorbidity (PRR 1.23, 95% CI 1.04–1.47). A CPP referral was less likely among patients with low education. The median diagnostic interval was 39 days (interquartile range: 15–72). When the GP initially did not suspect cancer, the likelihood of longer diagnostic interval increased.

Conclusion: The majority of patients with pancreatic cancer began their diagnostic route in general practice. Diagnosing pancreatic cancer swiftly in general practice was challenging; the GP did often not initially suspect cancer or refer to a CPP and several of the patient characteristics were associated with the GPs initial suspicion of cancer or CPP referral. Thus, there may be room for improvements in the diagnostics of pancreatic cancer in general practice.

KEY POINTS

- Patients with pancreatic cancer have a poor prognosis, as pancreatic cancer is often diagnosed in late stage.
- The majority of patients with pancreatic cancer began their diagnostic process in general practice.
- General practitioners (GPs) suspected cancer at the first consultation in one out of three patients with pancreatic cancer; more often in older and comorbid patients.
- The GPs suspicion of cancer was associated with urgent referral and shorter time to diagnosis.

ARTICLE HISTORY

Received 17 May 2021

Accepted 12 November 2021

KEYWORDS

Denmark; pancreatic neoplasms; early detection of cancer; general practice; cancer patient pathway

Background

Danish patients with pancreatic cancer have a poor prognosis with an estimated 5-year survival of less than 5% [1]. One explanation is that the majority of these patients are diagnosed with advanced disease stage and approximately 80% are not candidates for curative surgery [2]. To improve the prognosis in this patient group, more focus is needed on the possibilities to diagnose pancreatic cancer earlier in time [3].

As screening is not currently available for patients with pancreatic cancer [4], the first point of presentation is primarily through emergency presentation, incidental findings at hospitals in connection with investigation of other diseases, or GP referral (e.g. two-week-wait in the UK and Cancer Patient Pathways (CPP) in Denmark) [5–7]. Only few studies exist on the first point of presentation for patients with pancreatic cancer. An English study based on register data

CONTACT Line F. Virgilsen  line.virgilsen@ph.au.dk Research Unit for General Practice, Bartholins Allé 2, Aarhus C, 8000, Denmark

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

reported that 47% of patients were diagnosed after presentation in general practice [5], whereas two European studies based on survey data and interviews found that 70%–75% of the patients presented first in general practice [6,8]. One of these studies found that about half of these patients were referred to the two-week-wait system [8], which tends to reduce the time to diagnosis for most cancer types, including pancreatic cancer [9,10]. However, no data exist from a Nordic healthcare system.

Urgent referral to cancer diagnostics is often guided by symptoms [7]. Yet, pancreatic cancer is typically characterised by vague and unspecific symptoms [4,11,12]. This complicates the patients' healthcare seeking [8] and the GP's suspicion of cancer [9], which is a strong indicator for diagnostic action and shorter time to diagnosis in other cancer types [9]. So far, the evidence is sparse on the time intervals in the diagnostic journey for patients with pancreatic cancer [6,8,10,13,14], and little is known about the factors that might provide more rapid diagnosis of pancreatic cancer [6,8].

To obtain a faster and optimal diagnostic process hopefully leading to improvements in the survival of pancreatic cancer patients, more knowledge is needed about the diagnostic pathway. Therefore, the aim of this study was to investigate the first point of contact in patients diagnosed with pancreatic cancer, and to study factors associated with the GP's suspicion of cancer, CPP referral and long diagnostic interval.

Material and methods

Setting

This study was conducted in Denmark, which runs a tax-funded healthcare system with free and equal access for all citizens to most health services. A total of 98% of the population is registered with a general practice who acts as gatekeeper to the secondary health care system, except for emergencies, ear-nose-throat specialists and eye specialists who have direct access [15]. Since 2008/09, CPPs and standardised guidelines have been adopted to ensure equal, prompt and uniform diagnosis and treatment of most cancer types within specific time frames.

Design, study population and data collection

This study was conducted as a cross-sectional study and based on data from a series of six cohorts among newly diagnosed cancer patients in Denmark collected between 2005 and 2016. For this study, we included

patients from the cohort of 2010 and 2016 as these cohorts included data on pancreatic cancer patients and because all Danish Cancer Patient Pathways were implemented from 2010.

We included patients diagnosed with pancreatic cancer (International Classification of Diseases (ICD) Revision 10 code: C25 (except C254, endocrine pancreas) and C241, Ampulla of Vater) with no previous history of cancer. Patients from the two cohorts were identified in the Danish National Patient Register (DNPR) [16] in the period from 1 May to 31 August 2010 or from 1 July to 31 December 2016. The general practice of each included patient was identified in the Danish National Health Service Register [17]. The GP involved in the first presentation of symptoms, or who was most familiar with the patient, was asked to fill in the questionnaire based on their electronic patient record, to ensure that the responses were based on the observations made at the time of the first contact in the cancer course. Questionnaires were sent to the GPs of alive patients on a monthly basis in the data collection period to reduce the time span between the diagnosis and receipt of the questionnaire. However, in 2016, GP participation required written consent from the patient according to the Danish Health Act. For deceased patients in 2016, the Danish Patient Safety Authority granted permission for the GP to be contacted without patient consent so the GPs of these patients received the questionnaire 6–12 months after the patient was diagnosed with cancer. The GP questionnaires were developed by researchers at the Research Unit for General Practice in Aarhus, Denmark and the included time intervals followed the definition from the Aarhus Statement [18]. The questionnaires were pilot tested among GPs prior to data collection where minor changes were made to ensure each question were understood as intended.

Main variables

All data sources were linked at a personal level through the unique civil registration number assigned to all permanent Danish residents [19].

Four main variables were obtained from the GP questionnaire, and the responses were based on the GPs' clinical evaluation and information from their medical records. First, the first point of presentation was obtained from the question: "Based on your current knowledge, where did the patient present for the first time before the diagnosis?" Screening is not an option in this patient group, and none of the GPs responded that the patient was diagnosed through

specialised private practice or out-of-hours primary care. Thus, the possible routes included: presentation in general practice, diagnosed at the hospital after a contact for other disease, emergency presentation or other route. Second, GP's suspicion of cancer or serious illness was based on the question: "When the patient consulted general practice for the first time, what was your overall evaluation of the patient?" Three response options were: suspected cancer, suspected serious illness, did not suspect cancer or serious illness. Third, GP referral to a CPP after the first cancer-related presentation had the response options: yes and no. Fourth, the diagnostic interval was defined based on the Aarhus Statement [18] and assessed by calculating the interval between the date of the first presentation of cancer-related symptoms in general practice (as assessed by the GP and recorded in the medical record) and the date of diagnosis [18], which was obtained from the DNPR. The diagnostic interval was defined as the number of days between these two dates. Negative values were coded as 0 days, and intervals longer than one year were coded as 365 days in accordance with the procedures applied by the International Cancer Benchmarking Partnership (ICBP).

In addition, we obtained information on the patient's health and sociodemographic characteristics. Sex and age at diagnosis were obtained from the Danish Civil Registration System [19]. Statistics Denmark provided data on the patient's highest attained education, which was categorised according to the International Standard Classification of Education (ISCED) of the United Nations Education, Scientific and Cultural Organization (UNESCO) into short (≤ 10 years of education), medium (11–15 years of education) and long (> 15 years of education), and civil status was dichotomised into living alone or married/cohabiting. Charlson's Comorbidity Index (CCI) score was calculated from diagnoses (excluding cancer) recorded in the DNPR for up to ten years preceding the pancreatic cancer diagnosis, and CCI scores were categorised into none (score 0), low (scores 1–2) or high (scores > 2). Previous GP attendance was measured as the number of face-to-face consultations in general practice in the 12–24 months preceding the cancer diagnosis recorded in the Danish Health Service Register [20], and these were categorised based on the cut-off at the 50th and 75th percentiles, i.e. 0–1, 2–5 and > 5 consultations.

Statistical analyses

The point of first presentation was described by proportions and expressed by percentages. Generalised

linear models (GLMs) with log link for the Poisson family were used to study the association between patient characteristics and the likelihood of the GP to suspect cancer or serious illness. GLMs were also used to study patient characteristics, GP suspicion, the likelihood of CPP referral and the likelihood of long diagnostic interval. A long diagnostic interval was defined as the 75th percentile for the total population, i.e. 72 days. To comply with the data protection regulations (i.e. values less than 5 observations at the 75% cut-off are suppressed to prevent disclosure), a pseudo-variable for the diagnostic interval was generated using a method described elsewhere [21]. In all analyses, an unadjusted model was presented and followed by a model adjusted for sex, age, year of diagnosis and GP suspicion of cancer (except for GP suspicion in the analysis in which suspicion was the dependent variable).

Stata software, version 16.0, was used for all statistical analyses.

Results

In total, 438 patients were eligible for inclusion, and 69.2% ($n = 303$) of the GPs responded to the questionnaire.

Point of first presentation

The GPs provided information on the point of first presentation for 303 patients, and 259 (85.5%) of these patients presented with initial symptoms in general practice. The remaining 14.5% were diagnosed at the hospital after a contact related to another disease (6.9%), emergency presentation (5.6%) or other reasons (2.0%). Among the patients presenting to general practice, a higher proportion of patients were aged 70–79 years and had no comorbidity compared with patients presenting elsewhere (Table 1).

GP suspicion of cancer

Among the 259 patients who presented symptoms to their GP, the GP suspected cancer in 32.7%, other serious illness in 34.7%, and neither cancer nor other serious illness in 32.6% (Table 1). Male sex and older age were associated with higher likelihood of GP suspicion. Patients with high comorbidity were also more likely to raise suspicion in their GP compared with patients without comorbidity (PRR 1.23, 95% CI 1.04–1.47) (Table 2).

Table 1. Patient characteristics according to GP involvement ($n = 303$).

	GP involved		GP not involved		Total	
	<i>n</i>	(% column)	<i>n</i>	(% column)	<i>n</i>	(%)
Total	259	(100.0)	44	(100.0)	303	(100)
Age groups (years)						
<70	108	(41.7)	24	(54.5)	132	(43.6)
≥70–79	102	(39.4)	8	(18.2)	110	(36.3)
≥80	49	(18.9)	12	(27.3)	61	(20.1)
Sex						
Female	119	(45.9)	18	(40.9)	137	(45.2)
Male	140	(54.1)	26	(59.1)	166	(54.8)
Year of diagnosis						
2010	131	(50.6)	26	(59.1)	157	(51.8)
2016	128	(49.4)	18	(40.9)	146	(48.2)
Marital status						
Cohabiting/married	165	(63.7)	28	(63.6)	193	(63.7)
Living alone	94	(36.3)	16	(36.4)	110	(36.3)
Education						
Short	113	(43.6)	23	(52.3)	136	(44.9)
Medium and long ^a	146	(56.4)	21	(47.7)	167	(55.1)
Charlson's Comorbidity Index						
None (0)	159	(61.4)	18	(40.9)	177	(58.4)
Mild (1–2)	58	(22.4)	14	(31.8)	72	(23.8)
High (2 or more)	42	(16.2)	12	(27.3)	54	(17.8)
Previous use of primary care^b						
Infrequent attenders (0–1 visits)	60	(23.2)	8	(18.2)	68	(22.4)
Regular attenders (2–5 visits)	109	(42.1)	16	(36.4)	125	(41.3)
Frequent attenders (>5 visits)	90	(34.7)	20	(45.5)	110	(36.3)
Tumour stage						
Local/regional	69	(26.6)	9	(20.5)	78	(25.7)
Distant	119	(45.9)	17	(38.6)	136	(44.9)
Missing	71	(27.4)	18	(40.9)	89	(29.4)
Death within one year after diagnosis						
Not death	172	(66.4)	28	(63.6)	200	(66.0)
Death	87	(33.6)	16	(36.4)	103	(34.0)
GP suspicion at the first presentation						
Suspected cancer	82	(32.7)	NA	–	82	(32.7)
Suspected serious illness	87	(34.7)	NA	–	87	(34.7)
No suspicion	82	(32.6)	NA	–	82	(32.6)
Cancer Patient Pathway referral at the first GP presentation						
No	192	(77.1)	NA	–	192	(77.1)
Yes	57	(22.9)	NA	–	57	(22.9)

Significant differences between groups are shown in bold (Pearson's χ^2 square test).

Abbreviations: GP: general practitioner; NA: not applicable.

^aMedium and long education was combined because one of the columns had <5 observations, thus, cannot be reported according to Statistics Denmark's data protection regulations.

^bBased on the number of face-to-face consultations with the GP in the 12–24 months prior to diagnosis.

CPP referral by the GP

Among the 259 patients who presented symptoms to their GP, the GP referred 22.9% of the patients to a CPP at the first presentation (Table 1). Compared with less educated patients, patients with long education were more likely to be referred to a CPP at the first GP consultation (PRR 2.06, 95% CI 1.09–3.88) (Table 3). When the GP did not suspect cancer at the first consultation, the likelihood of CPP referral at the first presentation was reduced (PRR 0.46, 95% CI 0.25–0.85) (Table 3).

Diagnostic interval

The median diagnostic interval was 39 (Inter quartile interval: 15–72) days (median in 2010: 46 days, median

in 2016: 31 days) and 10% had a diagnostic interval longer than 173 days. When the GP did not suspect cancer, an almost three-fold higher likelihood of long diagnostic interval was observed (PRR 2.77, 95% CI 1.40–5.49) (Table 4). Patients who were referred to a CPP at the first GP consultation had a non-significant lower likelihood of long diagnostic interval (PRR 0.59, 95% CI 0.28–1.24). The patient's health and sociodemographic characteristics were not associated with the diagnostic interval (Table 4).

Discussion

Principal findings

This population-based study found that the majority of patients with pancreatic cancer began their route

Table 2. GP suspicion of cancer or serious illness at the first cancer-related presentation according to the patient's characteristics ($n = 251^a$).

	Prevalence rate ratio of GP suspecting cancer or serious illness					
	GP suspicious		Unadjusted		Adjusted ^b	
	N	% (row)	PRR	95% CI	PRR	95% CI
Total ($n = 251$, No GP suspicion, $n = 82$)	169	(67.3)	–	–	–	–
Sex						
Female ($n = 111$)	68	(61.2)	1	–	1	–
Male ($n = 140$)	101	(72.1)	1.18	(0.98–1.41)	1.20	(1.00–1.43)
Age groups (years)						
18–69 ($n = 104$)	57	(54.8)	1	–	1	–
70–79 ($n = 100$)	73	(73.0)	1.33	(1.08–1.65)	1.34	(1.09–1.66)
>80 ($n = 47$)	39	(83.0)	1.51	(1.22–1.88)	1.54	(1.24–1.91)
Year of diagnosis						
2010 ($n = 124$)	84	(67.7)	1	–	1	–
2016 ($n = 127$)	85	(66.9)	0.99	(0.83–1.17)	0.98	(0.82–1.15)
Marital status						
Cohabiting/married ($n = 161$)	107	(66.5)	1	–	1	–
Not married ($n = 90$)	62	(68.9)	1.04	(0.87–1.24)	1.03	(0.86–1.23)
Education						
Short ($n = 108$)	80	(74.1)	1	–	1	–
Medium ($n = 100$)	61	(61.0)	0.82	(0.68–1.00)	0.84	(0.69–1.02)
Long ($n = 43$)	28	(65.1)	0.88	(0.69–1.12)	0.94	(0.73–1.21)
Charlson's Comorbidity Index						
None (score 0) ($n = 154$)	100	(64.9)	1	–	1	–
Mild (score 1–2) ($n = 56$)	34	(60.7)	0.93	(0.73–1.19)	0.91	(0.72–1.15)
High (score 3 or more) ($n = 41$)	35	(85.4)	1.31	(1.11–1.56)	1.23	(1.04–1.47)
Previous use of primary care^c						
Infrequent attenders (0–1 visit) ($n = 56$)	38	(67.9)	1.03	(0.82–1.29)	1.13	(0.89–1.42)
Regular attenders (2–5 visits) ($n = 108$)	71	(65.7)	1	–	1	–
Frequent attenders (>5 visits) ($n = 87$)	60	(69.0)	1.05	(0.86–1.28)	1.02	(0.84–1.23)

GP: general practitioner; PRR: prevalence rate ratio; CI: confidence interval.

Significant results are shown in bold.

^a8 patients with missing information on GP's suspicion of cancer at the first cancer-related.

^bAdjusted for sex, age and year of diagnosis.

^cBased on number of face-to-face consultations with the GP in the 12–24 months prior to diagnosis and categorised based on the cut-off at the 25 and 75% percentiles.

to diagnosis in general practice. In one out of three patients, neither cancer nor serious illness was suspected when the patient first presented in general practice. GPs were more likely to suspect cancer in males, older patients and patients with high comorbidity. One in four patients were referred to a CPP after the first presentation in general practice, and CPP referral was more likely among highly educated patients. GP suspicion was crucial for both CPP referral at the first presentation and shorter diagnostic interval.

Strengths and limitations

Patients were identified through the population-based DNPR [22]. High data completeness ensured nearly complete identification and minimised selection bias. Questionnaire data was the primary source of data, and prior to data collection, the questionnaires were pilot tested among GPs to enhance the face and construct validity. The questionnaire focused on the first presentation in general practice and not the entire diagnostic process in general practice, which could

have provided additional insights for patients with several GP consultations before referral as often is the case with patients diagnosed with pancreatic cancer [23]. The response rate of 70% might have caused selection bias. However, the sub-analyses showed that patients' health and sociodemographic characteristics were equally distributed among responding and non-responding GPs, yet, a higher proportion of non-respondents (47%) were deceased within one year of the diagnosis compared to respondents (34%) (Data not shown).

Information bias caused by GP recall problems cannot be rejected. Yet, as GPs in Denmark have electronic medical records, which they were encouraged to consult, information bias from this source was minimised. The study included two cohorts of cancer patients diagnosed in 2010 and 2016. We have no systematically collected information on public campaigns in the time between the two cohorts, however, the data relied on GP's assessment which may be less susceptible to public campaigns. Further, all current CPPs were implemented from 2010 and the proportion of CPP referral and GP suspicion did not differ statistically

Table 3. Cancer Patient Pathway referral after the first cancer-related presentation according to the patient's characteristics ($n = 249$)^a.

	Prevalence rate ratio of any Cancer Patient Pathway referral at first presentation					
	Cancer Patient Pathway referral, yes		Unadjusted		Adjusted ^b	
	N	% (row)	PRR	95% CI	PRR	95% CI
Total ($n = 249$)	57	(22.9)	–	–	–	–
Sex						
Female ($n = 110$)	20	(18.2)	1	–	1	–
Male ($n = 139$)	37	(26.6)	1.46	(0.90–2.38)	1.44	(0.89–2.33)
Age groups (years)						
18–69 ($n = 104$)	18	(17.3)	1	–	1	–
70–79 ($n = 100$)	29	(29.0)	1.68	(0.99–2.82)	1.54	(0.91–2.60)
>80 ($n = 45$)	10	(22.2)	1.28	(0.64–2.56)	1.13	(0.55–2.30)
Year of diagnosis						
2010 ($n = 125$)	28	(22.4)	1	–	1	–
2016 ($n = 124$)	29	(23.4)	1.04	(0.66–1.65)	1.21	(0.77–1.91)
Marital status						
Cohabiting/married ($n = 158$)	37	(23.4)	1	–	1	–
Not married ($n = 91$)	20	(22.0)	0.94	(0.58–1.52)	1.13	(0.71–1.81)
Education						
Short ($n = 109$)	18	(16.5)	1	–	1	–
Medium ($n = 98$)	25	(25.5)	1.54	(0.90–2.66)	1.40	(0.79–2.49)
Long ($n = 42$)	14	(33.3)	2.02	(1.11–3.69)	2.06	(1.09–3.88)
Charlson's Comorbidity Index						
None (0) ($n = 151$)	36	(23.8)	1	–	1	–
Mild (1–2) ($n = 57$)	14	(24.6)	1.03	(0.60–1.76)	1.05	(0.63–1.73)
High (3 or more) ($n = 41$)	7	(17.1)	0.72	(0.34–1.49)	0.70	(0.33–1.48)
Previous use of primary care^c						
Infrequent attenders (0–1 visit) ($n = 56$)	15	(26.8)	1.33	(0.75–2.35)	1.35	(0.76–2.41)
Regular attenders (2–5 visits) ($n = 140$)	22	(15.7)	1	–	1	–
Frequent attenders (>5 visits) ($n = 53$)	20	(37.7)	1.18	(0.69–2.02)	1.21	(0.73–2.02)
GP suspicion at the first presentation						
Suspected cancer ($n = 80$)	31	(38.7)	1	–	1	–
Suspected serious illness ($n = 86$)	12	(14.0)	0.36	(0.20–0.65)	0.37	(0.20–0.69)
No suspicion ($n = 81$)	13	(16.0)	0.41	(0.23–0.73)	0.46	(0.25–0.85)

GP: general practitioner; PRR: prevalence rate ratio; CI: confidence interval.

Significant results are shown in bold.

^a10 patients with missing information on GP-initiated Cancer Patient Pathway referral after the first cancer-related presentation.

^bAdjusted for sex, age, year of diagnosis and GP's suspicion.

^cBased on face-to-face consultations with the GP in the 12–24 months prior to diagnosis and categorised based on the cut-off at the 25 and 75% percentiles.

significantly in the two cohorts (data not shown) indicating no major changes in GP behaviour over time.

Data on symptoms experienced by the patients at the first presentation could have provided additional insights into the mechanisms as symptoms may be linked with the point of first presentation, the GP's suspicion and the choice of action [9]. Unfortunately, these data was not available in this study as first presented symptoms were not collected in the GP questionnaire.

Due to the relatively small population size, the statistical precision was reduced, and type II errors cannot be ruled out. With a larger population, some of the findings may have reached statistical significance, e.g. the association between previous GP attendance and CPP referral. Finally, the implications for early diagnosis of pancreatic cancer apply only to the patients who presented in general practice and first time cancer patients, as patients with previous cancer were excluded to avoid that the GP suspicion, CPP referral

and diagnostic interval was modified by previous experiences.

Comparison with other studies

For the majority of the included patients, the first point of symptom presentation was in the GP setting. This finding is comparable with the findings in two European studies [6,8] which reported that the majority of patients initially presented in general practice, although the proportion of patients presenting as emergency was larger in both the UK [8] and France [6] than in our study.

The GP suspected cancer in one in three patients at the first presentation, which is lower than reported across all cancer types in Denmark, where the GP suspected cancer at the first presentation in 48% [9]. Still, it is roughly similar to the 27% reported in patients with lung cancer in Denmark [24]. This may reflect that pancreatic cancer often presents with unspecific

Table 4. Long diagnostic interval according to the patient's characteristics ($n = 223^a$).

	Prevalence rate ratio of long diagnostic interval (≥ 72 days)					
	Long diagnostic interval (>72 days)		Unadjusted		Adjusted ^b	
	N	% (row)	PRR	95% CI	PRR	95% CI
Total ($n = 223$)	56	(25.1)	–	–	–	–
Sex						
Female ($n = 102$)	29	(26.7)	1	–	1	–
Male ($n = 121$)	27	(23.8)	0.87	(0.55–1.37)	0.97	(0.62–1.51)
Age groups (years)						
18–69 ($n = 92$)	25	(27.4)	1	–	1	–
70–79 ($n = 93$)	22	(22.6)	0.84	(0.51–1.39)	0.93	(0.55–1.57)
>80 ($n = 38$)	9	(25.6)	0.94	(0.50–1.78)	1.17	(0.63–2.20)
Year of diagnosis						
2010 ($n = 119$)	30	(25.2)	1	–	1	–
2016 ($n = 104$)	26	(25.0)	1.01	(0.64–1.59)	0.97	(0.61–1.53)
Marital status						
Cohabiting/married ($n = 144$)	34	(23.6)	1	–	1	–
Not married ($n = 79$)	22	(27.8)	1.18	(0.74–1.87)	1.10	(0.69–1.76)
Education						
Short ($n = 100$)	27	(26.3)	1	–	1	–
Medium ($n = 87$)	18	(20.7)	0.79	(0.46–1.34)	0.80	(0.47–1.38)
Long ($n = 36$)	11	(32.4)	1.23	(0.70–2.19)	1.32	(0.76–2.31)
Charlson's Comorbidity Index						
None (score 0) ($n = 137$)	35	(26.3)	1	–	1	–
Mild (score 1–2) ($n = 50$)	14	(25.5)	0.96	(0.56–1.67)	0.90	(0.51–1.59)
High (score 3 or more) ($n = 36$)	7	(20.0)	0.73	(0.36–1.51)	0.80	(0.39–1.63)
Previous use of primary care^c						
Infrequent attenders (0–1 visit) ($n = 54$)	15	(26.3)	1.36	(0.75–2.45)	1.37	(0.74–2.53)
Regular attenders (2–5 visits) ($n = 94$)	21	(21.3)	1	–	1	–
Frequent attenders (>5 visits) ($n = 75$)	20	(27.6)	1.42	(0.83–2.42)	1.40	(0.84–2.33)
GP suspicion at the first presentation						
Suspected cancer ($n = 73$)	10	(13.7)	1	–	1	–
Suspected serious illness ($n = 74$)	17	(23.0)	1.75	(0.87–3.54)	1.75	(0.84–3.65)
No suspicion ($n = 72$)	27	(37.5)	2.74	(1.43–5.24)	2.77	(1.40–5.49)
Cancer Patient Pathway referral at the first GP presentation						
No ($n = 166$)	47	(28.3)	1	–	1	–
Yes ($n = 50$)	7	(14.0)	0.49	(0.24–1.03)	0.59	(0.28–1.24)

GP: general practitioner; PRR: prevalence rate ratio; CI: confidence interval.

Significant results are shown in bold.

^a36 patients with missing information on one of the dates used to calculate the diagnostic interval.

^bAdjusted for sex, age, year of diagnosis and GP's suspicion.

^cBased on number of face-to-face consultations with the GP in the 12–24 months prior to diagnosis and categorised based on the cut-off at the 25 and 75% percentiles.

and vague symptoms, supporting pancreatic cancer as a so-called 'hard to diagnose' type of cancer like lung cancer [23].

To our knowledge, the link between a patient's health and sociodemographic characteristics and the GPs' suspicion of cancer at the first presentation in general practice has not been studied previously among patients with pancreatic cancer. This study provides new knowledge, which indicates a higher propensity among GPs to suspect cancer or serious illness in older patients, in patients with high comorbidity and, to some extent, in males. In line with previous studies on other cancer types, we found that the GP's suspicion of cancer was strongly associated with a higher propensity of referring to a CPP after the patient's first clinical presentation and with shorter diagnostic interval [9,25]. This underlines that the GP's initial actions are important in healthcare systems with gatekeeping.

Initial CPP referral was observed in 22%, which was comparable to a previous study reporting that 27% of patients with lung cancer were referred to a CPP after the first GP presentation [24]. However, Walter et al. [8] reported that half of the patients with pancreatic cancer presenting to their GP in England were referred to a fast-track pathway. Walter et al. analysed the final referral mode and not the action by the GP at the first presentation. Thus, the different findings in the two studies could indicate that an interval may exist from the first presentation until urgent referral (e.g. CPP) for half of those referred to a fast-track pathway for pancreatic cancer [8].

An interesting finding in this study was that patients with low education were less likely to get a CPP referral at the first consultation, which is somewhat supported elsewhere [25]. Clinical bias has been raised as a potential explanation, as GPs may more actively refer affluent patients [26], or affluent patients

may be more successful in articulating and navigating the referral process in the cancer pathway [27]. However, the explanation for this finding remains unclear, and more research should be undertaken.

The diagnostic interval decreased from 2010 to 2016, which is in line with results in cancer patients in general [10,28], although the median diagnostic interval in this study was seven days longer than the diagnostic interval found for all cancer types combined in 2010 [28]. Nevertheless, the median length of the diagnostic interval of 39 days is similar to the diagnostic interval in English patients with pancreatic cancer [8], and it corresponds well with research indicating that patients with pancreatic cancer more often than patients with other types of cancer present three times or more in general practice before referral [23]. The patient's sociodemographic characteristics were not associated with the diagnostic interval, which could be due to the adjustment of cancer suspicion in the model, which seemed to be the factor with the largest impact on both CPP referral and diagnostic interval. Likewise, another study on pancreatic cancer patients found no association between sociodemographic characteristics and time intervals [8].

Implications

In countries with GP gatekeeping, the GP plays a crucial role in diagnosing cancer early. This study shows that additional support for the GPs is needed to ensure timely diagnosis of pancreatic cancer. Although a CPP for pancreatic cancer has been implemented in Denmark, three out of four patients with pancreatic cancers were not referred to a CPP at their first cancer-related consultation in general practice. As pancreatic cancer is difficult to diagnose due to vague and unspecific symptoms [2], studies into increased access to diagnostic tools, such as ultrasound, CT and endoscopy, is needed. Further, continuing medical education of GPs is crucial to ensure upgrading of the skills required for diagnosing pancreatic cancer. In light of the results, medical education could draw attention to how the educational level of the patient affect the GPs diagnostic behaviour as this study indicated lower propensity in referral of low educated patients.

Conclusion

The vast majority of patients with pancreatic cancer began their diagnostic route by presenting symptoms to their GP. The time to diagnosis was prolonged when the GP did not suspect pancreatic cancer.

Suspicion of cancer or other serious illness was associated with male gender, older age and high comorbidity. Finally, patients with low education were less often referred to a CPP.

Ethical approval

The project is registered in the Record of Processing Activities at the Research Unit for General Practice, Aarhus, in accordance with the Danish Data Protection Act (35) and the General Data Protection Regulation (GDPR) by the EU (36). As the data is based on registry and questionnaire data, approval by the Committee on Health Research Ethics in the Central Denmark Region was not required (37). Patients diagnosed in 2016 signed a consent allowing us to contact their GP and collect data on the diagnostic process of their patients. In accordance with section 46 of the Danish Health Act, the Danish Patient Safety Authority granted us permission to obtain questionnaire data based on the GPs' medical records for patients diagnosed in 2016 who deceased shortly after their diagnosis (case no. 3–3013-1956/1).

Disclosure statement

The authors declare that there are no conflicts of interests.

Funding

The study was supported by the M.L. Jørgensen and Gunner Hansen's Foundation, and the Danish General Practice Fund. The funders were not involved in any aspects of the study.

References

- [1] Cronin-Fenton DP, Erichsen R, Mortensen FV, et al. Pancreatic cancer survival in Central and Northern Denmark from 1998 through 2009: a population-based cohort study. *Clin Epidemiol*. 2011;3(Suppl 1): 19–25.
- [2] Kleeff J, Korc M, Apte M, et al. Pancreatic cancer. *Nat Rev Dis Primers*. 2016;2:16022.
- [3] Rasmussen LS, Fristrup CW, Jensen BV, et al. Initial treatment and survival in 4163 Danish patients with pancreatic cancer: a nationwide unselected real-world register study. *Eur J Cancer*. 2020;129:50–59.
- [4] Zhou B, Xu JW, Cheng YG, et al. Early detection of pancreatic cancer: where are we now and where are we going? *Int J Cancer*. 2017;141(2):231–241.
- [5] Elliss-Brookes L, McPhail S, Ives A, et al. Routes to diagnosis for cancer - determining the patient journey using multiple routine data sets. *Br J Cancer*. 2012; 107(8):1220–1226.
- [6] Jooste V, Dejardin O, Bouvier V, et al. Pancreatic cancer: wait times from presentation to treatment and survival in a population-based study. *Int J Cancer*. 2016;139(5):1073–1080.

- [7] Probst HB, Hussain ZB, Andersen O. Cancer patient pathways in Denmark as a joint effort between bureaucrats, health professionals and politicians—a national Danish project. *Health Policy*. 2012;105(1):65–70.
- [8] Walter FM, Mills K, Mendonça SC, et al. Symptoms and patient factors associated with diagnostic intervals for pancreatic cancer (SYMPTOM pancreatic study): a prospective cohort study. *The Lancet Gastroenterol Hepatol*. 2016;1(4):298–306.
- [9] Jensen H, Topping ML, Olesen F, et al. Cancer suspicion in general practice, urgent referral and time to diagnosis: a population-based GP survey and registry study. *BMC Cancer*. 2014;14:636.
- [10] Neal RD, Din NU, Hamilton W, et al. Comparison of cancer diagnostic intervals before and after implementation of NICE guidelines: analysis of data from the UK general practice research database. *Br J Cancer*. 2014;110(3):584–592.
- [11] Garg SK, Chari ST. Early detection of pancreatic cancer. *Curr Opin Gastroenterol*. 2020;36(5):456–461.
- [12] Singhi AD, Koay EJ, Chari ST, et al. Early detection of pancreatic cancer: opportunities and challenges. *Gastroenterology*. 2019;156(7):2024–2040.
- [13] Lyratzopoulos G, Saunders CL, Abel GA, et al. The relative length of the patient and the primary care interval in patients with 28 common and rarer cancers. *Br J Cancer*. 2015;112:S35–S40.
- [14] Keeble S, Abel GA, Saunders CL, et al. Variation in promptness of presentation among 10,297 patients subsequently diagnosed with one of 18 cancers: evidence from a national audit of cancer diagnosis in primary care. *Int J Cancer*. 2014;135(5):1220–1228.
- [15] Pedersen KM, Andersen JS, Sondergaard J. General practice and primary health care in Denmark. *J Am Board Fam Med*. 2012;25 (Suppl 1):S34–S8.
- [16] Lynge E, Sandegaard JL, Rebolj M. The Danish national patient register. *Scand J Public Health*. 2011;39(7 Suppl):30–33.
- [17] Andersen JS, Olivarius F, Krasnik A. The Danish national health service register. *Scand J Public Health*. 2011;39(7 Suppl):34–37.
- [18] Weller D, Vedsted P, Rubin G, et al. The Aarhus statement: improving design and reporting of studies on early cancer diagnosis. *Br J Cancer*. 2012;106(7):1262–1267.
- [19] Pedersen CB. The Danish civil registration system. *Scand J Public Health*. 2011;39(7 Suppl):22–25.
- [20] Olivarius NF, Hollnagel H, Krasnik A, et al. The Danish national health service register. A tool for primary health care research. *Dan Med Bull*. 1997;44(4):449–453.
- [21] StataHacks. Pseudo percentiles 2019. [Assessed the 04.10.21. Available from: http://w.bruunisejs.dk/StataHacks/Datamanagement/pseudo_percentiles/pseudo_percentiles/.
- [22] Larsen MB, Jensen H, Hansen RP, et al. Identification of patients with incident cancers using administrative registry data. *Dan Med J*. 2014;61(2):A4777.
- [23] Lyratzopoulos G, Neal RD, Barbiere JM, et al. Variation in number of general practitioner consultations before hospital referral for cancer: findings from the 2010 national cancer patient experience survey in England. *The Lancet Oncol*. 2012;13(4):353–365.
- [24] Guldbrandt LM, Fenger-Gron M, Rasmussen TR, et al. The role of general practice in routes to diagnosis of lung cancer in Denmark: a population-based study of general practice involvement, diagnostic activity and diagnostic intervals. *BMC Health Serv Res*. 2015;15:21.
- [25] Baun ML, Jensen H, Falborg AZ, et al. Ovarian cancer suspicion, urgent referral and time to diagnosis in Danish general practice: a population-based study. *Fam Pract*. 2019;36(6):751–757.
- [26] Forrest LF, Sowden S, Rubin G, et al. Socio-economic inequalities in stage at diagnosis, and in time intervals on the lung cancer pathway from first symptom to treatment: systematic review and Meta-analysis. *Thorax*. 2017;72(5):430–436.
- [27] Forrest LF, Adams J, White M, et al. Factors associated with timeliness of post-primary care referral, diagnosis and treatment for lung cancer: population-based, data-linkage study. *Br J Cancer*. 2014;111(9):1843–1851.
- [28] Jensen H, Topping ML, Olesen F, et al. Diagnostic intervals before and after implementation of cancer patient pathways - a GP survey and registry based comparison of three cohorts of cancer patients. *BMC Cancer*. 2015;15(1):308–015.