



Usual care and management of fall risk increasing drugs in older dizzy patients in Dutch general practice

Hanneke Stam, Thomas Harting, Marjolijn van der Sluijs, Rob van Marum, Henriëtte van der Horst, Johannes C. van der Wouden & Otto R. Maarsingh

To cite this article: Hanneke Stam, Thomas Harting, Marjolijn van der Sluijs, Rob van Marum, Henriëtte van der Horst, Johannes C. van der Wouden & Otto R. Maarsingh (2016) Usual care and management of fall risk increasing drugs in older dizzy patients in Dutch general practice, *Scandinavian Journal of Primary Health Care*, 34:2, 165-171, DOI: [10.3109/02813432.2016.1160634](https://doi.org/10.3109/02813432.2016.1160634)

To link to this article: <https://doi.org/10.3109/02813432.2016.1160634>



© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 06 Apr 2016.



Submit your article to this journal [↗](#)



Article views: 1289



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)



RESEARCH ARTICLE



Usual care and management of fall risk increasing drugs in older dizzy patients in Dutch general practice

Hanneke Stam^a, Thomas Harting^a, Marjolijn van der Sluijs^a, Rob van Marum^{a,b}, Henriëtte van der Horst^a, Johannes C. van der Wouden^a and Otto R. Maarsingh^a

^aDepartment of General Practice and Elderly Care Medicine and EMGO Institute for Health and Care Research, VU University Medical Center, Amsterdam, The Netherlands; ^bDepartment of Geriatric Medicine, Jeroen Bosch Hospital, 's-Hertogenbosch, The Netherlands

ABSTRACT

Objective: For general practitioners (GPs) dizziness is a challenging condition to deal with. Data on the management of dizziness in older patients are mostly lacking. Furthermore, it is unknown whether GPs attempt to decrease Fall Risk Increasing Drugs (FRIDs) use in the management of dizziness in older patients. The aim of this study is to gain more insight into GP's management of dizziness in older patients, including FRID evaluation and adjustment. **Design:** Data were derived from electronic medical records, obtained over a 12-month period in 2013. **Setting:** Forty-six Dutch general practices. **Patients:** The study sample comprised of 2812 older dizzy patients of 65 years and over. Patients were identified using International Classification of Primary Care codes and free text. **Main outcome measures:** Usual care was categorized into wait-and-see strategy (no treatment initiated); education and advice; additional testing; medication adjustment; and referral. **Results:** Frequently applied treatments included a wait-and-see strategy (28.4%) and education and advice (28.0%). Additional testing was performed in 26.8%; 19.0% of the patients were referred. Of the patients 87.2% had at least one FRID prescription. During the observation period, GPs adjusted the use of one or more FRIDs for 11.7% of the patients. **Conclusion:** This study revealed a wide variety in management strategies for dizziness in older adults. The referral rate for dizziness was high compared to prior research. Although many older dizzy patients use at least one FRID, FRID evaluation and adjustment is scarce. We expect that more FRID adjustments may reduce dizziness and dizziness-related impairment.

KEY POINTS

- It is important to know how general practitioners manage dizziness in older patients in order to assess potential cues for improvement.
- This study revealed a wide variety in management strategies for dizziness in older patients.
- There was a scarcity in Fall Risk Increasing Drug (FRID) evaluation and adjustment.
- The referral rate for dizziness was high compared with previous research.

ARTICLE HISTORY

Received 27 October 2015
Accepted 11 February 2016

KEYWORDS

Aged; dizziness; drug prescriptions; general practice; therapy; The Netherlands

Introduction

For general practitioners (GPs) dizziness is a challenging condition to deal with: dizziness may refer to a variety of sensations and the complaint dizziness may accompany harmless but also very serious conditions. Dizziness can refer to several sensations including a giddy or rotational sensation, a loss of balance, a faint feeling, light-headedness, instability or unsteadiness, a tendency to fall, or a feeling of everything turning black.[1] There is a broad etiologic spectrum of peripheral, central (neurological), and general medical causes

for dizziness. Furthermore, several authors suggest that dizziness in older people might be a multifactorial geriatric syndrome.[2–6] A geriatric syndrome is defined as a specific symptom that is caused by multiple underlying factors, involving multiple organ systems that tend to contribute to the geriatric syndrome.[7]

It is important to know how GPs manage dizziness in older patients, in order to assess potential cues for improvement. Only few studies focused on the management of dizziness in primary care but these studies did not focus on older patients.[8–12]

CONTACT Hanneke Stam, MD ✉ stam.h@vumc.nl VU University Medical Center, Department of General Practice and Elderly Care Medicine and EMGO Institute for Health and Care Research, Van der Boechorststraat 7, 1081 BT Amsterdam, The Netherlands

© 2016 The Author(s). Published by Taylor & Francis. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

A medication review should be part of the assessment of older patients with dizziness because it is assumed that medication is a contributory factor to dizziness in as much as 25% of these patients.[13] However, it is unknown whether GPs consider if medication might contribute to the dizziness when they evaluate an older patient. Drugs that contribute to dizziness [14,15] demonstrate a striking similarity with the list of Fall Risk Increasing Drugs (FRIDs).[16] This similarity might be explained by the fact that dizziness increases the risk of falling [17] and FRIDs are known to affect postural control.[18] Therefore, the list of FRIDs might be a useful proxy for potential dizziness inducing medication. Recently, Harun and Agrawal recommended to reconsider the use of FRIDs when evaluating and treating dizzy patients.[19]

The aim of this study is to gain more insight into the management of dizziness in older adults in general practice, with a focus on FRID evaluation and adjustment.

Material and methods

We used anonymized data from the database of the Academic Network of General Practice of VU University Medical Center (ANH-VUmc). The ANH-VUmc is a collaboration between VU University Medical Center and general practices located in an urban area of the Netherlands. The ANH-VUmc database contains anonymized routine health care data. Observational studies based on the ANH-VUmc database are carried out according to Dutch privacy legislation and are exempted from informed consent of patients.

Identification of patients

An electronic search strategy was applied to identify our target population: all patients aged 65 years and above who visited their GP because of dizziness in 2013. The database was searched for International Classification of Primary Care (ICPC) codes N17 "vertigo/dizziness" and H82 "vertiginous syndrome". Additionally, we searched for dizziness in the full text records by searching for the Dutch equivalents of "dizz*" and "vertigo".

We extracted the following data from anonymized patient records: patient characteristics (gender, age), characteristics of consultation for dizziness (type of consultation, date, ICPC diagnosis given by the GPs), characteristics of prescribed drugs (Anatomical Therapeutic Chemical (ATC) classification, prescription date) and information on symptoms, physical examination, diagnostic evaluation, and treatment.

Reviewing electronic full-text medical records

Two authors, a medical doctor (M.S.) and a medical student (T.H.), manually reviewed the electronic full-text consultations of the identified patients. They both reviewed and discussed a randomly selected 10% of the data to increase the reliability of the data extraction.

During review, patients and consultations were excluded for several reasons: (1) patients with a consultation that mentioned dizziness when dizziness was not an actual problem for the patient were excluded (e.g. "not dizzy"); (2) if a GP was consulted through a third party (e.g. a family member); and (3) only the first three consultations for dizziness during the observation period were coded to reduce the influence of patients with many consultations.

In case of multiple consultations, the ICPC code of the second or third consultation was considered the final diagnosis. If the GP recorded two ICPC codes in the chronologically last consultation, we recorded both of the ICPC codes as final diagnosis. Treatment was categorized as follows: wait-and-see strategy (no treatment initiated); education and advice; additional testing; medication adjustment; and referral. All treatment modalities were categorized, including multiple treatment modalities in one consultation and in multiple consultations (e.g. a wait-and-see strategy and blood analysis in the first consultation and referral in the second consultation in one patient was all recorded). Category medication adjustment was only used in case of a dose reduction of FRIDs, discontinuation of FRIDs, and/or the prescription of antivertigo drugs and antiemetics. The list of FRIDs included psychotropic drugs (sedatives, antidepressants, neuroleptics), cardiovascular drugs (antihypertensives, nitrates, antiarrhythmic agents, nicotinic acid, β -adrenoceptor blocker eye drops) and other drugs (analgesics, antivertigo drugs, hypoglycemics, urinary antispasmodics).[16]

Data analysis

We used descriptive analyses to describe the study population and to categorize treatment modalities. Practice list sizes were available and prevalence rates were calculated using the mid-time population. Age groups were compared with Chi-square tests and logistic regression analysis.

Results

Figure 1 presents a flowchart of the study selection process. A total of 2812 older dizzy patients were included in the sample.




Figure 1. Flowchart of study selection process of 2812 older dizzy patients. GP: General Practitioner; EMR: Electronic Medical Record.

Table 1. Overview of characteristics of 2812 older dizzy patients.

Characteristics	n (%)
Gender, female	1892 (67.3)
Age in years, mean (range)	77.0 (65–101)
65–74	1180 (42.0)
75–84	1086 (38.6)
≥85	546 (19.4)
Diagnosis ^a	
Symptom diagnoses	901 (32.0)
Cardiovascular condition	513 (18.2)
Peripheral vestibular disease	294 (10.5)
Infections	135 (4.8)
Psychiatric condition	73 (2.6)
Musculoskeletal condition	42 (1.5)
Neurological condition ^b	37 (1.3)
Metabolic or endocrine condition	32 (1.1)
Adverse effect medical agents	29 (1.0)
Other	668 (23.8)
No diagnosis recorded	110 (3.9)

^aDoes not add up to 100% because the GPs recorded two ICPC codes for 22 patients.

^bexcl. cerebrovascular conditions.

Table 1 presents the clinical features of the total study population of 2812 older dizzy patients. The median age of the population was 76 years (range: 65–101). The majority of the patients was female (67.3%). The 12-month prevalence of dizziness was 11.8%. The prevalence of dizziness significantly increased with age (χ^2 (linear-by-linear) = 354, df = 1, $p < 0.001$). The median consultation frequency for dizziness was 1 (range: 1–23), 444 patients

(15.8%) had more than 3 consultations in the dizziness episode. The mean follow-up time after the first consultation was 199 days (range: 1–365), 94.7% of the sample could be followed up more than one month. The most frequently recorded diagnoses for dizziness were symptom diagnoses (32.0%), cardiovascular conditions (18.2%), and peripheral vestibular diseases (10.5%).

Table 2 provides an overview of treatment modalities. Frequently applied treatments by GPs were a strategy of wait-and-see ($n = 799$, 28.4%) and providing education and advice ($n = 786$, 28.0%). Additional tests were performed to 755 patients (26.8%), of which blood analyses ($n = 622$, 22.1%) were most often carried out. Medication was prescribed and adjusted in 526 patients (18.7%). Finally, 533 patients (19.0%) were referred to a medical specialist. Patients were most often referred to a neurologist ($n = 136$, 4.8%), cardiologist ($n = 110$, 3.9%), and physiotherapist ($n = 65$, 2.3%).

The use of FRIDs and the frequencies of all FRID medication adjustments are displayed in Table 3. The patients were prescribed a mean of 3.1 FRIDs (SD 2.1). As many as 87.2% of the patients had at least one FRID prescription. FRID's were adjusted in 330 patients (11.7%). GPs reduced a FRID dose for 111 patients (3.9%) and discontinued FRID for 199 patients (7.1%). For 20 patients GPs both reduced the FRID dose and discontinued a FRID. Dose reductions of FRIDs

Table 2. Overview of management of 2812 older dizzy patients.

	n (%)
Wait-and-see (no treatment)	
Total	799 (28.4)
Education and advice	
Vestibular training exercises	87 (3.10)
Breathing exercises	5 (0.20)
Other education or advice	709 (25.2)
Total ^a	786 (28.0)
Additional test	
Blood analysis	622 (22.1)
Urine analysis	89 (3.20)
Electrocardiography	65 (2.30)
24-h blood pressure monitoring	41 (1.50)
Other	34 (1.20)
Total ^a	755 (26.8)
Medication prescription and medication adjustment	
Prescription of antiemetics	83 (3.00)
Prescription of antivertigo drugs	143 (5.10)
in dizziness caused by Ménière's disease	7 (0.20)
in other dizziness of vestibular origin	52 (1.80)
in other types of dizziness	84 (3.00)
Adjustment of FRIDs	330 (11.7)
dose reduction	131 (4.70)
discontinuation	219 (7.80)
Total ^a	526 (18.7)
Referral	
Neurologist	136 (4.80)
Cardiologist	110 (3.90)
Physical therapist	65 (2.30)
Internist	58 (2.10)
Otolaryngologist	37 (1.30)
Geriatrician	25 (0.90)
Ophthalmologist	19 (0.70)
Psychotherapist	16 (0.60)
Other	112 (4.00)
Total ^a	533 (19.0)

^aTotal number of patients. As some patients had multiple tests, adjustments or referrals, the sum of individual items does not add up to the total.

BPPV: benign paroxysmal positional vertigo; FRIDs: fall risk increasing drugs.

significantly increased with age (10 year odds ratio 1.36; 95% confidence interval 1.10–1.69).

Discussion

Principal findings

We performed this study to gain more insight into the management of dizziness in older adults in general practice. Frequent treatments included a wait-and-see strategy (28.4%) and education and advice (28.0%). Additional tests were carried out for 26.8% of the sample. For 11.7% of the patients GPs adjusted FRID prescription. The frequency of dose reductions of FRIDs significantly increased with age. GPs referred 19.0% of the older dizzy patients to specialized care.

The prevalence of dizziness was 11.8%. Cardiovascular conditions and peripheral vestibular disease were most often recorded as cause of dizziness. The GP recorded a symptom diagnosis in 32% of the patients. In 3.9% of the patients the GP did not record a diagnosis at all.

Strengths and weaknesses

Prior to this study, few studies have investigated the management of dizziness in older patients in general practice.[8–10] To our knowledge, this is the first study that investigates adjustments of FRIDs for older dizzy patients.

In the Netherlands, all patients are registered with a GP. The GP provides care and acts as a gatekeeper to specialized care. As a consequence, data presented in this study are a proper reflection of the prevalence and management of older dizzy people in general practice.

By using a dataset derived from electronic medical records (EMRs) we were able to identify a large sample of older patients with dizziness. However, we note that the quality of data depends on the accuracy of registration by GPs. Yet, GPs who participate in the general practice-based registration network where data for this study is derived from, are annually trained on registering and coding of medical data.

Findings in relation to other studies

Only a small number of studies investigated the management of dizziness in general practice.[8–11] A wait-and-see strategy was also frequently seen in a previous study with dizzy patients of both younger and older age.[9] Observation, reassurance, and advice to change behavior tended to be used in older dizzy patients in this study of Sloane et al. [9] In three studies with younger and older adults, drugs were prescribed to 60–90% of the patients,[8–10] which is much more frequent than in our sample. The high rate of referral to specialized care (19.0%) in this study is remarkable; international studies reported 4–16% referrals [8–10] and Dutch studies reported referral rates of 3.2–4.5%.[20,21] Several studies demonstrate that GPs' referral decisions are influenced by a complex mix of patient, physician, and health care system structural characteristics.[22–24] As the Dutch health care system did not change, changes in patient's expectations, doctor's perceptions of patients expectations reassurance for the patient) might have influenced the referral rate. It is unknown whether the referrals were effective in achieving their objectives and whether they were cost-effective.

This is the first study that focussed on the use and adjustments of FRIDs in older dizzy patients. FRID use was quite high in our study sample, with a mean of 3.1 FRID prescriptions per patient. This is similar to the mean of FRID prescriptions in a study of older patients

Table 3. Use, adjustments, and new prescription of FRIDs of 2812 older dizzy patients.

Drug group	Use of FRIDs <i>n</i> (%) ^a	Dose reductions <i>n</i> (%) ^b	Discontinuation <i>n</i> (%) ^b	Newly prescribed <i>n</i> (%) ^b
Cardiovascular FRIDs				
Diuretics	1199 (42.6)	27 (2.3)	56 (4.7)	
β-Blockers	1144 (40.7)	35 (3.1)	24 (2.1)	
Calcium channel blockers	654 (23.3)	19 (2.9)	33 (5.0)	
Angiotensin converting enzyme inhibitors	714 (25.4)	17 (2.4)	19 (2.7)	
Angiotensin receptor blockers	620 (22.0)	13 (2.1)	9 (1.5)	
Nitrates	343 (12.2)	3 (0.9)	4 (1.2)	
Antiarrhythmic agents	61 (2.22)	1 (1.6)	0 (0)	
Digoxin	97 (3.42)	0 (0)	0 (0)	
Psychotropic FRIDs				
Antivertigo drugs	239 (8.52)	0 (0)	7 (2.9)	143 (59.8)
Analgesics (opioids)	524 (18.6)	13 (2.5)	23 (4.4)	
Anxiolytics and hypnotics	815 (29.0)	5 (0.6)	8 (1.0)	
Antidepressants	389 (13.8)	4 (1.0)	8 (2.1)	
Neuroleptics	91 (3.22)	2 (2.2)	3 (3.3)	
α-blockers and anticholinergics	330 (11.7)	1 (0.3)	14 (4.2)	
Hypoglycaemics	508 (18.1)	3 (0.6)	3 (0.6)	
Antihistamines	292 (10.4)	1 (0.3)	1 (0.3)	
β-Blocker eye drops	98 (3.52)	0 (0)	0 (0)	
Other FRIDs	471 (16.7)	3 (0.6)	27 (0.6)	
No FRID use	359 (12.8)	na	na	na
Total medication adjustments	na	131 (4.7) ^c	219 (7.8) ^d	(140) ^e

^aPercent of patients being prescribed at least one drug of the displayed fall risk increasing drug categories during 2013.

^bPercent of patients with at least one FRID adjustment per FRID group.

^cIn 14 patients, two medications were reduced in dose, in one patient three medications were reduced in dose.

^dIn 20 patients two medications were stopped.

^eIn three patients two antivertigo drugs were prescribed.

FRID: Fall Risk Increasing Drug; na: not applicable.

plus a fall history and a study with frail aged patients (3.3 and 3.4, respectively).[16,25]

GPs discontinued a FRID or reduced the dose of a FRID for 11.7% of the patients. Other management strategies were carried out in 19.0–28.4% of our sample. Compared with other management strategies, FRID adjustment was carried out the least frequent. In a qualitative study on situations in which GPs associate FRIDs with falls, drug use was often not perceived as a prominent factor.[26] One situation leading to a consideration of the drug prescribed was if a patient had fallen or presented with a symptom such as dizziness. However, the paradox of not being able to predict the outcome of changes in drug treatment was perceived as challenging and uncomfortable; the GPs believed that it might be better not to change prescriptions instead.[26] In other research on de-prescribing of medication, four main themes of doctor-related barriers are described: lack of awareness on consequences of polypharmacy; inertia or devolving of responsibility; lack of skills and knowledge; and presumed lack of feasibility.[27] Evidently, adjustment of medication seems difficult. However, given the fact that 87.2% of the patients were having at least one prescribed FRID in this sample, there is ample room for improvement by adjusting more FRID medication. A medication review, and evaluation and adjustment of FRIDs in particular, may be a simple and effective management strategy to reduce dizziness and dizziness-related

impairment in older patients. FRIDs can also be adjusted if the cause of dizziness has not yet been identified.

The prevalence of dizziness in older adults seems higher than previously reported. Maarsingh et al. reported a dizziness prevalence of 8.3%, whereas this study revealed a dizziness prevalence of 11.8% in a highly comparable sample.[28] Furthermore, Sloane et al. reported a dizziness prevalence of 7.0% in patients aged 85 years and above.[8] Drug prescription has increased,[29] with higher rates of adverse drug reactions as a result. This may have resulted in a higher prevalence of dizziness in older patients, as adverse drug reactions are thought to contribute to dizziness.[13] On the other hand, GPs who participate in the general practice-based registration network where data for the current study is derived from, are annually trained on registering and coding of medical data. This may have caused higher registration rates by GPs in this study. It is important to continue monitoring whether dizziness prevalence in older adults is rising, because this will increase the burden of dizziness on society, health care systems, and individuals.

In 35.9% of this sample GPs recorded a symptom diagnosis or did not record a diagnosis at all. In a similar study, GPs recorded a symptom diagnosis in 40.0% of the patients.[28] This high rate of unknown cause of dizziness may be the result of difficulties for GPs to establish the origin of dizziness.

Meaning of the study: implications for clinicians and research

Compared with other management strategies, FRID adjustments were carried out the least. We recommend to always evaluate the use of FRIDs for older dizzy patients and to consider adjustment. The use of FRIDs should be discontinued if no health risks are involved by discontinuation or the dose of FRIDs should be reduced when discontinuation is not an option.

The referral rate for dizziness was high compared with previous research. Therefore, it is important to investigate whether referrals for dizziness are effective in achieving their objectives and whether they are cost-effective.

Acknowledgements

The authors wish to thank J.H.K. Joosten for her assistance with the data extraction. Furthermore the authors would like to thank all the GPs who participate in the Academic Network of General Practitioners of the VU University Medical Center.

Ethical approval

The ANH-VUmc database contains anonymized routine health care data. Observational studies based on the ANH-VUmc database are carried out according to Dutch privacy legislation and are exempted from informed consent of patients.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the article.

Funding information

This study was funded by the Netherlands Organisation for Health Research and Development. The funding body did not and will not have any role in the design, collection, analysis, or interpretation of data, nor in the writing of the manuscript.

References

- [1] Sloane PD, Coeytaux RR, Beck RS, et al. Dizziness: state of the science. *Ann Intern Med.* 2001;134(Pt 2): 823–832.
- [2] Kao AC, Nanda A, Williams CS, et al. Validation of dizziness as a possible geriatric syndrome. *J Am Geriatr Soc.* 2001;49:72–75.
- [3] Gassmann KG, Rupprecht R. Dizziness in an older community dwelling population: a multifactorial syndrome. *J Nutr Health Aging.* 2009;13:278–282.
- [4] Gomez F, Curcio CL, Duque G. Dizziness as a geriatric condition among rural community-dwelling older adults. *J Nutr Health Aging.* 2011;15:490–497.
- [5] De Moraes SA, Soares WJ, Ferrioli E, et al. Prevalence and correlates of dizziness in community-dwelling older people: a cross sectional population based study. *BMC Geriatr.* 2013;13.
- [6] Maarsingh OR, Stam H, van de Ven PM, et al. Predictors of dizziness in older persons: a 10-year prospective cohort study in the community. *BMC Geriatr.* 2014;14:133.
- [7] Inouye SK, Studenski S, Tinetti ME, et al. Geriatric syndromes: clinical, research, and policy implications of a core geriatric concept. *J Am Geriatr Soc.* 2007;55: 780–791.
- [8] Sloane PD. Dizziness in primary care. Results from the National Ambulatory Medical Care Survey. *J Fam Pract.* 1989;29:33–38.
- [9] Sloane PD, Dallara J, Roach C, et al. Management of dizziness in primary care. *J Am Board Fam Med.* 1994; 7:1–8.
- [10] Bird JC, Beynon GJ, Prevost AT, et al. An analysis of referral patterns for dizziness in the primary care setting. *Br J Gen Pract.* 1998;48:1828–1832.
- [11] Jayarajan V, Rajenderkumar D. A survey of dizziness management in General Practice. *J Laryngol Otol.* 2003;117:599–604.
- [12] Nazareth I, Yardley L, Owen N, et al. Outcome of symptoms of dizziness in a general practice community sample. *Fam Pract.* 1999;16:616–618.
- [13] Maarsingh OR, Dros J, Schellevis FG, et al. Causes of persistent dizziness in elderly patients in primary care. *Ann Fam Med.* 2010;8:196–205.
- [14] Lin E, Aligene K. Pharmacology of balance and dizziness. *NeuroRehabilitation.* 2013;32:529–542.
- [15] Shoaib OA, Nyandeghe AN, Slattum PW. Medication-related dizziness in the older adult. *Otolaryngol Clin North Am.* 2011;44:455–471.
- [16] Van der Velde N, Stricker BHC, Pols HA, et al. Risk of falls after withdrawal of fall-risk-increasing drugs: a prospective cohort study. *Br J Clin Pharmacol.* 2007;63: 232–237.
- [17] Tinetti ME, Doucette J, Claus E, et al. Risk factors for serious injury during falls by older persons in the community. *J Am Geriatr Soc.* 1995;43:1214–1221.
- [18] De Groot MH, van Campen JPCM, Moek MA, et al. The effects of fall-risk-increasing drugs on postural control: a literature review. *Drug Aging.* 2013;30:901–920.
- [19] Harun A, Agrawal Y. The use of Fall Risk Increasing Drugs (FRIDs) in patients with dizziness presenting to a neurotology clinic. *Otol Neurotol.* 2015;36:862–864.
- [20] Okkes IM, Oskam SK, Lamberts H. Van klacht naar diagnose. Episodegegevens uit de huisartspraktijk [From complaint to diagnosis. Episode data from family practice]. Couthinho: Bussum; 1998.
- [21] Cardol M, van Dijk L, de Jong JD, et al. Tweede Nationale Studie naar ziekten en verrichtingen in de huisartspraktijk. Huisartsenzorg: wat doet de poortwachter? [Second National Study into diseases and actions in general practice: complaints and disorders

- in the population and in general practice]. Utrecht/Bilthoven: Nivel/RIVM; 2004.
- [22] Little P, Dorward M, Warner G, et al. Importance of patient pressure and perceived pressure and perceived medical need for investigations, referral, and prescribing in primary care: nested observational study. *BMJ*. 2004;328:444.
 - [23] Forrest CB, Nutting PA, von Schrader S, et al. Primary care physician specialty referral decision making: patient, physician, and health care system determinants. *Med Decis Mak*. 2006;26:76–85.
 - [24] Ringberg U, Fleten N, Førde OH. Examining the variation in GPs' referral practice: a cross-sectional study of GPs' reasons for referral. *Br J Gen Pract*. 2014;64: e426–e433.
 - [25] Bennett DA, Gnjdjic D, Gillett M, et al. Prevalence and impact of fall-risk-increasing drugs, polypharmacy, and drug-drug interactions in robust versus frail hospitalised falls patients: a prospective cohort study. *Drug Aging*. 2014;31:225–232.
 - [26] Bell HT, Steinsbekk A, Granas AG. Factors influencing prescribing of fall-risk-increasing drugs to the elderly: a qualitative study. *Scand J Prim Health Care*. 2015;33:107–114.
 - [27] Anderson K, Stowasser D, Freeman C, et al. Prescriber barriers and enablers to minimising potentially inappropriate medications in adults: a systematic review and thematic synthesis. *BMJ Open*. 2014;4:e006544.
 - [28] Maarsingh OR, Dros J, Schellevis FG, et al. Dizziness reported by elderly patients in family practice: prevalence, incidence, and clinical characteristics. *BMC Fam Pract*. 2010;11.
 - [29] Health, United States. With special feature on prescription drugs. U.S. Department of Health and Humans: Hyattsville, 2013; [cited 2015 Jun 12]. Available from: <http://www.cdc.gov/nchs/data/hus/hus13.pdf>.