

## DNA methylation profiles and biomarkers of oral squamous cell carcinoma

Yu-Fen Li, Yi-Hsiu Hsiao, Yi-Hui Lai, Yi-Chen Chen, Ying-Ju Chen, Jian-Liang Chou, Michael W Y Chan, Yu-Hsing Lin, Yung-An Tsou, Ming-Hsui Tsai & Chien-Kuo Tai

**To cite this article:** Yu-Fen Li, Yi-Hsiu Hsiao, Yi-Hui Lai, Yi-Chen Chen, Ying-Ju Chen, Jian-Liang Chou, Michael W Y Chan, Yu-Hsing Lin, Yung-An Tsou, Ming-Hsui Tsai & Chien-Kuo Tai (2015) DNA methylation profiles and biomarkers of oral squamous cell carcinoma , Epigenetics, 10:3, 229-236, DOI: [10.1080/15592294.2015.1006506](https://doi.org/10.1080/15592294.2015.1006506)

**To link to this article:** <https://doi.org/10.1080/15592294.2015.1006506>



Published online: 20 Mar 2015.



Submit your article to this journal [↗](#)



Article views: 2055



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 16 View citing articles [↗](#)

# DNA methylation profiles and biomarkers of oral squamous cell carcinoma

Yu-Fen Li<sup>1</sup>, Yi-Hsiu Hsiao<sup>1</sup>, Yi-Hui Lai<sup>2</sup>, Yi-Chen Chen<sup>2</sup>, Ying-Ju Chen<sup>2</sup>, Jian-Liang Chou<sup>2</sup>, Michael W Y Chan<sup>2</sup>, Yu-Hsing Lin<sup>2,3</sup>, Yung-An Tsou<sup>4</sup>, Ming-Hsui Tsai<sup>4</sup>, and Chien-Kuo Tai<sup>2,\*</sup>

<sup>1</sup>Institute of Biostatistics; China Medical University; Taichung, Taiwan; <sup>2</sup>Department of Life Science and Institutes of Molecular Biology and Biomedical Science; National Chung Cheng University; Min-Hsiung, Chia-Yi, Taiwan; <sup>3</sup>Department of Otolaryngology; Buddhist Dalin Tzu Chi General Hospital; Chia-Yi, Taiwan; <sup>4</sup>Department of Otolaryngology; China Medical University Hospital; Taichung, Taiwan

**Keywords:** DNA methylation array, oral squamous cell carcinoma, biomarkers, pyrosequencing, *FLT4*, *KDR*, *TFPI2*

Oral squamous cell carcinoma (OSCC) constitutes >90% of oral cancers and is the sixth most common malignancy among males worldwide and the fourth leading cause of death due to cancer among males in Taiwan. However, most patients do not receive a diagnosis of OSCC until the late stages, which have a lower survival rate. The use of molecular marker analysis to identify early-stage OSCC would permit optimal timing for treatments and consequently prolong survival. The aim of this study was to identify biomarkers of OSCC using the Illumina GoldenGate Methylation Cancer Panel, which comprised a total of 1,505 CpG sites covering 807 genes. Samples of buccal mucosa resected from 40 OSCC patients and normal tissue samples obtained from 15 patients (normal mucosa from OSCC patients or from patients undergoing surgery unrelated to OSCC) were analyzed. Fms-related tyrosine kinase 4 (*FLT4*) methylation exhibited a perfect specificity for detecting OSCC, with an area under the receiver operating characteristic curve of 0.91 for both all-stage and early-stage OSCC. Methylation of 7 genes (*ASCL1*, *FGF3*, *FLT4*, *GAS7*, *KDR*, *TERT*, and *TFPI2*) constitutes the top-20 panels for detecting OSCC. The top-20 panels for detecting early-stage OSCC contain 8 genes: *ADCYAP1*, *EPHA7*, *FLT4*, *GSTM2*, *KDR*, *MT1A*, *NPY*, and *TFPI2*. *FLT4* RNA expression and methylation level were validated using RT-PCR and a pyrosequencing methylation assay. The median level of *FLT4* expression was 2.14-fold for normal relative to OSCC tissue samples ( $P < 0.0001$ ). Among the 8 pyrosequenced *FLT4* CpG sites, methylation level was much higher in the OSCC samples. In conclusion, methylation statuses of selected genes, and especially *FLT4*, *KDR*, and *TFPI2*, might be of great potential as biomarkers for early detection of buccal OSCC.

## Introduction

Oral cancer is the sixth most common malignancy among males worldwide in terms of both incidence and mortality.<sup>1</sup> Particularly, oral cancer has been the fourth leading cause of death due to cancer among males in Taiwan since 2003.<sup>2</sup> The median age at death from oral cancer in 2010 was 56 years in men, which was the youngest age at death among the top-10 cancers in men. Moreover, oral cancer has had the highest rank in incidence and mortality rate among men aged 25–44 years in Taiwan since 2005.<sup>2</sup>

Oral squamous cell carcinoma (OSCC) constitutes more than 90% of oral cancers arising from the oral cavity.<sup>3</sup> The treatment of OSCC with early detection has a good outcome, but the 5-year survival rate is <30% among patients with stage IV disease.<sup>4</sup> Despite the ease of access to the oral cavity for clinical examination, OSCC is usually only diagnosed at advanced stages due to the patient's lack of awareness of any major problems.<sup>5</sup> The development of a convenient tool for detecting OSCC at an early stage would not only improve the survival of patients, but also reduce the associated medical expenses. This makes the search for biomarkers for early detection of OSCC an urgent

priority. The common risk factors for OSCC in most countries are cigarette smoking and alcohol drinking, and certain biomarkers for OSCC due to these risk factors have been identified; however, the main risk factor for OSCC in Taiwan is chewing of betel-quid, and the biomarkers for OSCC due to betel-quid consumption may be quite different from those identified for other risk factors.<sup>6,7</sup>

Epigenetic modifications, including DNA methylation and numerous histone modifications, are responsible for the altered gene expression patterns that allow for specific phenotypes.<sup>8</sup> DNA methylation plays an important role in normal mammalian development, but aberrant methylation patterns are correlated with several differentiation-related diseases, including various types of human cancers. Investigation of the promoter methylation of tumor-suppressor genes in the setting of oral epithelial dysplasia seems appropriate, given the relatively high frequency of this epigenetic change in OSCC.<sup>9</sup> Early epigenetic changes could predispose cells to further genetic abnormalities that may allow progression of the neoplastic process. Thus, identification of gene methylation as an appropriate marker may provide a sensitive method for OSCC detection.

\*Correspondence to: Chien-Kuo Tai; Email: biockt@ccu.edu.tw

Submitted: 11/12/2014; Revised: 12/20/2014; Accepted: 01/03/2015

<http://dx.doi.org/10.1080/15592294.2015.1006506>

The development of new technologies, such as DNA microarray analysis and next-generation sequencing, has enabled the expansion of DNA-methylation studies beyond just several candidate genes. Several methods of determining the occurrence of promoter methylation have been described in the literature, including methylation-specific PCR (MSP), combined bisulfite restriction analysis, bisulfite sequencing assay, pyrosequencing methylation assay, and microarray-based methylation analysis; the overwhelming majority of published data were obtained using the MSP method. Literature reviews<sup>10-12</sup> have revealed a wide variation in the reported sensitivity and specificity for this method, ranging from 30% to 90%. Given the limited number of CpG sites detected by MSP and the inconsistent results obtained using this method, we developed a diagnostic tool for OSCC using a high-throughput, quantitative methylation platform—the Illumina GoldenGate Methylation Array—to allow a more comprehensive profiling of DNA methylation.

## Results

### Sample characteristics for the Illumina GoldenGate Methylation Array

Selected sample characteristics are presented in Table 1. The mean age at surgery of the 40 OSCC cases from whom tissue was obtained for analysis was 52.6 years (range: 36–73 years), while the mean age of the subjects in the normal control group was 47.2 years (range: 17–70 years). Of the 40 OSCC samples, 40% were obtained from patients who were regarded as having early-stage OSCC: 22.5% at stage I and 17.5% at stage II.

### Identification of single CpG sites as biomarkers for OSCC

Among 1,505 CpG sites screened in the array, 34 CpG sites covering 29 genes exhibited a reasonable ability to distinguish OSCC (n=40) and normal (n=15) samples, with a specificity and sensitivity of  $\geq 70\%$  (Table 2). Moreover, the specificity of *FLT4*, *TFPI2*, *ASCL1*, and *KDR* was 100%, while the sensitivities of these genes for detecting OSCC were 83%, 78%, 75%, and 70%, respectively. The areas under the receiving operating characteristic curves (AUCs) decreased after leave-one-out cross validation, leaving only *FLT4* with an AUC of  $>0.8$ . The heat map of hierarchical cluster analysis for the 34 CpG sites is

**Table 1.** Characteristics of OSCC patients and normal specimens in the Illumina GoldenGate Methylation Array analysis

Characteristics	OSCC group (n=40)		Normal group* (n=15)	
Age (years, mean $\pm$ SE)	52.6 $\pm$ 9.6		47.2 $\pm$ 14.6	
Pathological Stage	n	%	n	%
Normal	—	—	5	33.3%
I	9	22.5%	8	53.3%
II	7	17.5%	—	—
III	10	25.0%	2	13.3%
IV	14	35.0%	—	—

SE: standard error.

\*10 samples are the normal tissues adjacent to OSCC tissues.

**Table 2.** Performance of a single CpG site for detecting OSCC

Rank <sup>†</sup>	CpG site	Discrimination statistics				Cross validation*	
		SPE	SEN	AUC	(95% CI)	AUC	(95% CI)
1	FLT4_E206_F	1.00	0.83	0.91	(0.85, 0.97)	0.83	(0.71, 0.94)
2	TFPI2_P9_F	1.00	0.78	0.89	(0.82, 0.95)	0.78	(0.64, 0.91)
3	ASCL1_E24_F	1.00	0.75	0.88	(0.81, 0.94)	0.75	(0.61, 0.89)
4	ADCYAP1_P455_R	0.87	0.88	0.87	(0.77, 0.97)	0.76	(0.58, 0.94)
5	MT1A_P49_R	0.87	0.85	0.86	(0.75, 0.96)	0.74	(0.56, 0.92)
7	ESR1_P151_R	0.80	0.90	0.85	(0.74, 0.96)	0.72	(0.52, 0.92)
7	KDR_E79_F	1.00	0.70	0.85	(0.78, 0.92)	0.70	(0.56, 0.84)
7	MYOD1_E156_F	0.80	0.90	0.85	(0.74, 0.96)	0.72	(0.52, 0.92)
10	EPHA5_E158_R	0.87	0.83	0.85	(0.74, 0.95)	0.72	(0.54, 0.89)
10	NPY_E31_R	0.87	0.83	0.85	(0.74, 0.95)	0.72	(0.54, 0.89)
10	RASGRF1_E16_F	0.87	0.83	0.85	(0.74, 0.95)	0.72	(0.54, 0.89)
12.5	GAS7_E148_F	0.93	0.75	0.84	(0.75, 0.94)	0.70	(0.54, 0.86)
12.5	TERT_E20_F	0.93	0.75	0.84	(0.75, 0.94)	0.70	(0.54, 0.86)
15	EPHA7_E6_F	0.87	0.80	0.83	(0.72, 0.94)	0.69	(0.51, 0.87)
15	HOXA5_P1324_F	0.87	0.80	0.83	(0.72, 0.94)	0.69	(0.51, 0.87)
15	NTRK3_P752_F	0.87	0.80	0.83	(0.72, 0.94)	0.69	(0.51, 0.87)
17.5	EPHA7_P205_R	0.87	0.78	0.82	(0.71, 0.93)	0.67	(0.49, 0.85)
17.5	MYH11_P22_F	0.87	0.78	0.82	(0.71, 0.93)	0.67	(0.49, 0.85)
19.5	FGF3_P171_R	0.93	0.70	0.82	(0.72, 0.91)	0.65	(0.49, 0.82)
19.5	HS3ST2_P171_F	0.73	0.90	0.82	(0.69, 0.94)	0.66	(0.44, 0.88)
21.5	GABRB3_E42_F	0.80	0.83	0.81	(0.69, 0.93)	0.66	(0.46, 0.86)
21.5	TFPI2_P152_R	0.80	0.83	0.81	(0.69, 0.93)	0.66	(0.46, 0.86)
23.5	BMP3_E147_F	0.87	0.75	0.81	(0.70, 0.92)	0.65	(0.47, 0.83)
23.5	GPX3_E178_F	0.87	0.75	0.81	(0.70, 0.92)	0.65	(0.47, 0.83)
25	PENK_P447_R	0.73	0.88	0.80	(0.68, 0.93)	0.64	(0.43, 0.86)
26	HOXB13_P17_R	0.80	0.80	0.80	(0.68, 0.92)	0.64	(0.44, 0.84)
27.5	HS3ST2_P546_F	0.73	0.85	0.79	(0.66, 0.92)	0.62	(0.41, 0.84)
27.5	TERT_P360_R	0.73	0.85	0.79	(0.66, 0.92)	0.62	(0.41, 0.84)
29	CHGA_E52_F	0.80	0.78	0.79	(0.66, 0.91)	0.62	(0.43, 0.81)
30	FRZB_E186_R	0.87	0.70	0.78	(0.67, 0.90)	0.61	(0.43, 0.78)
31	IHH_E186_F	0.80	0.75	0.78	(0.65, 0.90)	0.60	(0.41, 0.79)
32.5	ISL1_E87_R	0.73	0.75	0.74	(0.61, 0.88)	0.55	(0.35, 0.75)
32.5	ISL1_P379_F	0.73	0.75	0.74	(0.61, 0.88)	0.55	(0.35, 0.75)
34	MME_P388_F	0.73	0.73	0.73	(0.59, 0.86)	0.53	(0.33, 0.73)

SPE=sensitivity; SEN=sensitivity; AUC=area under the ROC curve; CI=confidence interval.

<sup>†</sup>ranked by AUC (an average rank is assigned for tied AUCs).

\*leave-one-out cross validation.

presented in Figure 1A. Of the 3 clusters, the right one comprised only OSCC samples, while most of the samples classified in the other 2 clusters were OSCC and normal samples. This implies that these 34 CpG sites are informative for detecting OSCC.

A subset analysis was conducted for the ability to detect early-stage OSCC (stages I and II). Among the 17 CpG sites covering 15 genes that had both a specificity and sensitivity of  $>70\%$  to discriminate the early-stage OSCC (n=16) and normal samples (n=15), the top-3 informative CpG sites (*FLT4\_E206*, *KDR\_E79*, and *TFPI2\_P9*) had a specificity of 100% and a sensitivity of  $\geq 75\%$  (Table 3). The AUCs decreased after leave-one-out cross validation, again leaving only *FLT4*, with an AUC of  $>0.8$ . The heat map presented in Figure 1B for distinguishing early-stage OSCC from normal samples shows 3 clusters (from left to right): (1) a mixture of

**Table 3.** Performance of a single CpG site for detecting early-stage OSCC

Rank <sup>†</sup>	CpG site	Discrimination statistics				Cross validation*	
		SPE	SEN	AUC	(95% CI)	AUC	(95% CI)
1	FLT4_E206_F	1.00	0.81	0.91	(0.81, 1.00)	0.81	(0.62, 1.00)
2.5	KDR_E79_F	1.00	0.75	0.88	(0.77, 0.98)	0.75	(0.53, 0.97)
2.5	TFPI2_P9_F	1.00	0.75	0.88	(0.77, 0.98)	0.75	(0.53, 0.97)
5	GSTM2_E153_F	0.87	0.81	0.81	(0.67, 0.95)	0.65	(0.42, 0.88)
5	MT1A_P49_R	0.87	0.81	0.84	(0.71, 0.97)	0.70	(0.48, 0.93)
5	NPY_E31_R	0.87	0.81	0.84	(0.71, 0.97)	0.70	(0.48, 0.93)
8.5	ADCYAP1_P455_R	0.87	0.75	0.81	(0.67, 0.95)	0.65	(0.42, 0.88)
8.5	EPHA7_E6_F	0.87	0.75	0.81	(0.67, 0.95)	0.65	(0.42, 0.88)
8.5	ESR1_P151_R	0.80	0.88	0.84	(0.70, 0.97)	0.70	(0.47, 0.93)
8.5	MYOD1_E156_F	0.80	0.88	0.84	(0.70, 0.97)	0.70	(0.47, 0.93)
11.5	HOXB13_P17_R	0.80	0.81	0.81	(0.66, 0.95)	0.65	(0.42, 0.88)
11.5	TFPI2_P152_R	0.80	0.81	0.81	(0.66, 0.95)	0.65	(0.42, 0.88)
13.5	HS3ST2_P171_F	0.73	0.88	0.80	(0.66, 0.95)	0.64	(0.40, 0.88)
13.5	PENK_P447_R	0.73	0.88	0.80	(0.66, 0.95)	0.64	(0.40, 0.88)
15	GABRB3_E42_F	0.80	0.75	0.78	(0.62, 0.93)	0.60	(0.36, 0.84)
16.5	HS3ST2_P546_F	0.73	0.75	0.74	(0.58, 0.90)	0.55	(0.31, 0.79)
16.5	TERT_P360_R	0.73	0.75	0.74	(0.58, 0.90)	0.55	(0.31, 0.79)

SPE=sensitivity ; SEN=sensitivity ; AUC=area under the ROC curve; CI=confidence interval.

†ranked by AUC (an average rank is assigned for tied AUCs).

\*leave-one-out cross validation.

OSCC and normal samples; (2) outnumbered by normal samples; and (3) OSCC samples only.

The hypermethylation status of the CpG sites in the Illumina GoldenGate Methylation Cancer Panel was found to be informative for detecting OSCC, while the hypomethylation status of the CpG sites was less informative (data not shown). The hypomethylation status of the studied CpG sites exhibited a good specificity but a poor sensitivity, and *ERN1* was the only gene with an AUC of >0.70 (AUC=0.72, specificity=93%, and sensitivity=50%).

### Identification of CpG panels as biomarkers for OSCC

Since there are numerous combinations for CpG panels, only the top-20 panels are listed in Tables 4 and 5. The top-20 CpG panels for detecting OSCC contain 7 genes: *ASCL1*, *FGF3*, *FLT4*, *GAS7*, *KDR*, *TERT*, and *TFPI2* (Table 4). The best panel was the combination of *FLT4* and *ASCL1*, with a specificity, sensitivity, and AUC of 100%, 90%, and 0.95, respectively. Adding *ASCL1* to the panel increased the sensitivity from 82.5% to 90% for *FLT4* alone (100% specificity). The best panel was followed by another 3 panels with the same performance, but with one extra CpG site that was considered redundant.

For detecting early-stage OSCC, the top-20 panels comprised 8 genes: *ADCYAP1*, *EPHA7*, *FLT4*, *GSTM2*, *KDR*, *MT1A*, *NPY*, and *TFPI2* (Table 5). Three panels (*KDR*+*GSTM2*+*MT1A*, *FLT4*+*GSTM2*+*MT1A*, and *GSTM2*+*MT1A*+*NPY*) exhibited the same dissemination ability (specificity=86.7%, sensitivity=100%, and AUC=0.933), followed by *FLT4* alone (specificity=100%, sensitivity=81.3%, and AUC=0.906).

### Validation of *FLT4* methylation by pyrosequencing assay

The methylation status of *FLT4* appeared to be a promising candidate as a biomarker for detecting OSCC. In fact, *FLT4\_E206* was the top biomarker and after leave-one-out cross validation was the only CpG site with an AUC of >0.80 for detecting all-stage or early-stage OSCC (highest rank in Tables 2 and 3). We validated this finding both internally and externally using the pyrosequencing methylation assay. Among those subjected to Illumina GoldenGate Methylation Array analysis, 9 normal samples and 17 OSCC samples with available DNA left were used for internal validation. Another 28 OSCC tissue specimens obtained from Buddhist Tzu Chi General Hospital served as samples for external validation. Among the 8 pyrosequenced CpG sites (+204, +206, +210, +225, +228, +237, +239, and +247) of *FLT4*, the differences in methylation levels between normal and OSCC samples were all statistically significant in the internal comparison (Fig. 2A and B). Six (+204, +206, +225, +228, +237, and +239) of the 8 pyrosequenced CpG sites exhibited statistically significant differences between normal and OSCC samples in the external comparison (Fig. 2C). As expected, the methylation levels were much higher in the OSCC samples (median methylation percentage range for the 8 CpG sites, 46–68%) than in the normal samples (median methylation percentage range, 10–23%).

### Validation of *FLT4* expression


Comparison of *FLT4* RNA expression between 21 paired normal and OSCC tissue samples yielded results that were in line with those obtained using the Illumina GoldenGate Methylation Array and pyrosequencing assays. The median level of *FLT4* expression in normal samples relative to OSCC samples was 2.14-fold (interquartile range=3.21;  $P<0.0001$ ) (Fig. 3).

## Discussion

Using an array approach, we observed that the methylation of CpG sites had both high sensitivity and specificity as biomarkers for detecting buccal OSCC. The hypermethylation of *FLT4* had perfect specificity for detecting all-stage or early-stage OSCC, and a sensitivity and AUC of >0.80 and >0.90, respectively. In addition, using the selected CpG combination panels slightly improved the ability to detect OSCC.

Based on the selected CpG panels, we identified 12 genes (*ADCYAP1*, *ASCL1*, *EPHA7*, *FGF3*, *FLT4*, *GAS7*, *GSTM2*, *KDR*, *MT1A*, *NPY*, *TERT*, and *TFPI2*) with great potential for detecting OSCC. The functions of the selected genes strengthen their potential role as biomarkers. *EPHA7*,<sup>13</sup> *FGF3*,<sup>14</sup> and *NPY*<sup>15,16</sup> are related to tumor progression, *GAS7* controls neurogenesis,<sup>17</sup> *ASCL1* plays a role in the early stages of development of specific neural lineages,<sup>18</sup> *TERT* regulates telomerase activity,<sup>19</sup> *ADCYAP1* is associated with hormone activity,<sup>20</sup> *GSTM2* belongs to a well-known phase II detoxifying enzyme family,<sup>21</sup> *MT1A* regulates cell growth and differentiation,<sup>22</sup> *TFPI-2* counteracts the activity of several extracellular matrix-associated proteases<sup>23,24</sup> and suppresses tumor invasion and metastasis,<sup>25-27</sup> and





**Figure 1.** Hierarchical clustering heat maps and dendrograms for distinguishing (A) OSCC from normal samples and (B) early-stage OSCC from normal samples. Mean  $\beta$  values from the selected CpG sites were used for hierarchical agglomerative clustering based on the Manhattan distance and complete linkage. Low, medium, and high methylation levels are indicated in green, black, and red, respectively.

FLT4 and KDR are receptors for vascular endothelial growth factors.<sup>28-31</sup> It has been reported that hypermethylation of *EPHA7*,<sup>32,33</sup> *GSTM2*,<sup>34</sup> *MT1A*,<sup>35</sup> *NPY*,<sup>36</sup> and *TERT*<sup>36-38</sup> are associated with cancers other than OSCC.

Viet et al.<sup>10</sup> used the Illumina GoldenGate Methylation Array to identify potential biomarkers for oral cancer by comparing the saliva samples of oral cancer patients before and after resection of their tumors. About one half of the identified genes (e.g., *ADCYAP1*, *FGF3*, *GAS7*, and *KDR*) also exhibited potential in our tissue samples as biomarkers of oral cancer. In a recent study, Jithesh et al.<sup>39</sup> also employed the same array and suggested that *EPHA5*, *GSTM2*, *HS3ST2*, *MME*, *MT1A*, *MYOD1*, *NPY*, and *PENK* are potential biomarkers for detecting oral cancer, as in the present study. The only triple-hit gene among the studies of Viet et al.,<sup>10</sup> Jithesh et al.,<sup>39</sup> and ourselves was *EPHA5*. In addition, Nagata et al.<sup>40</sup> reported that the aberrant methylation of a combination of *ECAD*, *TMEFF2*, *RAR $\beta$* , and *MGMT* from rinse samples can detect OSCC with a sensitivity and specificity of >90% by using noninvasive methods to retrieve samples and a different methylation platform. However, with the exception of *TMEFF2*, the results were not consistent with those of the present study. These inconsistencies can be explained by several factors, including differences in the sample types analyzed (tissue versus saliva/oral rinse) and different mechanisms of tumorigenesis (betel-quid chewing vs. cigarette smoking/alcohol drinking). Furthermore, the influence of race/ethnicity, tumor heterogeneity, experimental design, the methylation array platform employed, and the smallness of the samples in the studies cannot be excluded.

**Table 4.** Performance of a panel of CpG sites for detecting OSCC

Rank <sup>†</sup>	Panel of CpG sites	Specificity	Sensitivity	AUC
1	FLT4_E206_F + ASCL1_E24_F	100.0%	90.0%	0.950
3	FLT4_E206_F + ASCL1_E24_F + KDR_E79_F	100.0%	90.0%	0.950
3	TFPI2_P9_F + ASCL1_E24_F + KDR_E79_F	100.0%	90.0%	0.950
3	FLT4_E206_F + TFPI2_P9_F + ASCL1_E24_F	100.0%	90.0%	0.950
5.5	FLT4_E206_F + KDR_E79_F	100.0%	87.5%	0.938
5.5	TFPI2_P9_F + ASCL1_E24_F	100.0%	87.5%	0.938
7	FLT4_E206_F + TFPI2_P9_F + KDR_E79_F	100.0%	87.5%	0.938
8	FLT4_E206_F + TFPI2_P9_F	100.0%	85.0%	0.920
13	FLT4_E206_F + ASCL1_E24_F + TERT_E20_F	93.3%	90.0%	0.917
13	FLT4_E206_F + ASCL1_E24_F + FGF3_P171_R	93.3%	90.0%	0.917
13	ASCL1_E24_F + KDR_E79_F + TERT_E20_F	93.3%	90.0%	0.917
13	ASCL1_E24_F + KDR_E79_F + FGF3_P171_R	93.3%	90.0%	0.917
13	FLT4_E206_F + ASCL1_E24_F + GAS7_E148_F	93.3%	90.0%	0.917
13	FLT4_E206_F + TFPI2_P9_F + GAS7_E148_F	93.3%	90.0%	0.917
13	TFPI2_P9_F + ASCL1_E24_F + FGF3_P171_R	93.3%	90.0%	0.917
13	TFPI2_P9_F + KDR_E79_F + GAS7_E148_F	93.3%	90.0%	0.917
13	FLT4_E206_F + KDR_E79_F + GAS7_E148_F	93.3%	90.0%	0.917
18	FLT4_E206_F	100.0%	82.5%	0.913
19.5	TFPI2_P9_F + KDR_E79_F	100.0%	82.5%	0.913
19.5	ASCL1_E24_F + KDR_E79_F	100.0%	82.5%	0.913

<sup>†</sup>ranked by AUC and then the number of CpG sites in each combination (average rank is assigned if tied).


The epigenetic signature associated with human papillomavirus (HPV)-driven head and neck squamous cell carcinoma (HNSCC) has been described recently,<sup>41,42</sup> and it appears that HPV-16 might be a good prognostic biomarker for oropharyngeal cancer.<sup>43</sup> The status of HPV infection was not available for

the present samples, and so it was not possible to evaluate the influence of HPV on the performance of biomarkers. However, it has been shown that HPV does not play an important role in the tumorigenesis of betel-quid-related oral cancers.<sup>44</sup> Another limitation of the present study is that personal information such


**Table 5.** Performance of a panel of CpG sites for detecting early-stage OSCC

Rank <sup>†</sup>	Panel of CpG sites	Specificity	Sensitivity	AUC
2	KDR_E79_F + GSTM2_E153_F + MT1A_P49_R	86.7%	100.0%	0.933
2	FLT4_E206_F + GSTM2_E153_F + MT1A_P49_R	86.7%	100.0%	0.933
2	GSTM2_E153_F + MT1A_P49_R + NPY_E31_R	86.7%	100.0%	0.933
4	FLT4_E206_F	100.0%	81.3%	0.906
6	KDR_E79_F + TFPI2_P9_F	100.0%	81.3%	0.906
6	FLT4_E206_F + KDR_E79_F	100.0%	81.3%	0.906
6	FLT4_E206_F + TFPI2_P9_F	100.0%	81.3%	0.906
8	FLT4_E206_F + KDR_E79_F + TFPI2_P9_F	100.0%	81.3%	0.906
10.5	GSTM2_E153_F + NPY_E31_R	86.7%	93.8%	0.902
10.5	KDR_E79_F + GSTM2_E153_F	86.7%	93.8%	0.902
10.5	FLT4_E206_F + GSTM2_E153_F	86.7%	93.8%	0.902
10.5	GSTM2_E153_F + MT1A_P49_R	86.7%	93.8%	0.902
20	FLT4_E206_F + GSTM2_E153_F + EPHA7_E6_F	86.7%	93.8%	0.902
20	FLT4_E206_F + GSTM2_E153_F + ADCYAP1_P455_R	86.7%	93.8%	0.902
20	FLT4_E206_F + TFPI2_P9_F + GSTM2_E153_F	86.7%	93.8%	0.902
20	TFPI2_P9_F + GSTM2_E153_F + MT1A_P49_R	86.7%	93.8%	0.902
20	KDR_E79_F + TFPI2_P9_F + GSTM2_E153_F	86.7%	93.8%	0.902
20	GSTM2_E153_F + MT1A_P49_R + EPHA7_E6_F	86.7%	93.8%	0.902
20	GSTM2_E153_F + NPY_E31_R + EPHA7_E6_F	86.7%	93.8%	0.902
20	KDR_E79_F + GSTM2_E153_F + NPY_E31_R	86.7%	93.8%	0.902
20	FLT4_E206_F + GSTM2_E153_F + NPY_E31_R	86.7%	93.8%	0.902
20	FLT4_E206_F + KDR_E79_F + GSTM2_E153_F	86.7%	93.8%	0.902
20	GSTM2_E153_F + NPY_E31_R + ADCYAP1_P455_R	86.7%	93.8%	0.902
20	KDR_E79_F + GSTM2_E153_F + ADCYAP1_P455_R	86.7%	93.8%	0.902
20	KDR_E79_F + GSTM2_E153_F + EPHA7_E6_F	86.7%	93.8%	0.902
20	GSTM2_E153_F + MT1A_P49_R + ADCYAP1_P455_R	86.7%	93.8%	0.902
20	TFPI2_P9_F + GSTM2_E153_F + NPY_E31_R	86.7%	93.8%	0.902

<sup>†</sup>ranked by AUC and then the number of CpG sites in each combination (average rank is assigned if tied).



**Figure 2.** Box plots of methylation levels of 8 *FLT4* CpG sites determined by pyrosequencing methylation assay. (A) Normal samples. (B) OSCC samples for internal validation. (C) OSCC samples for external validation. In each box plot, the whiskers represent the 10th and 90th percentiles, the lower and upper limits of the box indicate the 25th and 75th percentiles. An asterisk mark (\*) indicates that the difference between normal and tumor samples reaches statistical significance.



**Figure 3.** Comparison of *FLT4* RNA expression between 21 paired normal and OSCC tissue samples. The mRNA level of *FLT4* in each normal tissue sample was compared with that of its corresponding OSCC tissue sample, which was assigned a value of 1. Error bars indicate standard deviations.

as exposures to tobacco, alcohol, and betel-quid was not available. Therefore, we cannot assess the influence of smoking, alcohol drinking, and betel-quid chewing in the analysis of the association between OSCC and DNA methylation.

The perfect specificity and high sensitivity of the biomarkers observed in the present study is very promising. However, the application of biomarkers to disease screening or detection relies not only on their accuracy, but also on their convenience. The tissue samples used in the present study were obtained using an invasive method, which is generally considered inconvenient for fast cancer screening. Gene methylation in saliva has been tested for the follow-up and early detection of still-curable relapses in HNSCC patients.<sup>45</sup> In addition, the methylation biomarkers selected in the present study should be examined for other oral cancer sites such as tongue, with a view to their further application.

In conclusion, methylation statuses of selected genes, and especially *FLT4*, *KDR*, and *TFPI2*, might be of great potential as biomarkers for the early detection of buccal OSCC. These candidate biomarkers provide a prospective application for screening and diagnosis; however, further studies are needed to evaluate samples obtained using noninvasive methods.

## Materials and Methods

### Illumina GoldenGate Methylation Array

Eight hundred and 7 cancer-related genes (1,505 CpG sites) were profiled in the Illumina GoldenGate Methylation Cancer Panel<sup>36</sup> by applying bisulfite sequencing technology on a microarray platform. The background normalization algorithm is used to minimize background variation within the microarray by using built-in negative control signals. The  $\beta$  value is used to estimate the methylation level of the CpG locus using the ratio of intensities between methylated and unmethylated alleles.<sup>36</sup>

### Tissue specimens for the methylation array profiling

Oral specimens were obtained from the tissue bank of China Medical University Hospital, Taiwan. The study was approved by the Ethics Committee of each participating hospital. Oral tissues from male individuals were collected and assigned to either the case group or the normal control group. The case group comprised resected tumor tissue harvested from 40 OSCC patients.

The control group comprised tissue samples harvested from 15 patients; in 10 cases the normal tissue was taken from an area near the tumor lesions in the oral cavities of OSCC patients in the case group, and in 5 cases the normal tissues were taken from the oral cavities of patients undergoing surgery unrelated to OSCC.

#### Tissue specimens for pyrosequencing methylation assay and qRT-PCR

A subset of 26 samples (9 normal and 17 OSCC tissues) among those oral specimens that were subjected to the Illumina GoldenGate Methylation Array (Illumina, Inc.) were also pyrosequenced. Another 28 fresh OSCC tissues that were obtained from Buddhist Tzu Chi General Hospital served as the external validation samples; 21 of these OSCC tissue samples accompanying their corresponding non-tumor oral tissues were used for measuring RNA expression level using qRT-PCR.

#### Pyrosequencing methylation assay

Genomic DNA of tissues was isolated using Gentra Puregene Tissue Kit (Qiagen) and 500 ng of each genomic DNA was bisulfite-modified using EZ DNA methylation kit (Zymo Research). Pyrosequencing primers were designed for the region of interest using Pyromark Assay Design v2.0 (Qiagen): forward, 5'-TTGGTAGGGGAGGGGGTAT-3'; reverse, 5'-biotin-CCCCCCCCCTTCCTCATCCC-3'; sequencing primer, 5'-GGATAAGGTGGTTTTGGT-3'. Pyrosequencing assay was carried out using the PyroMark Gold Q96 Reagents (Qiagen) in PyroMark Q24 System (Qiagen) according to the manufacturer's instructions.

#### Quantitative RT-PCR

Total RNA was extracted from fresh tissues using TRIzol reagent (Invitrogen) according to the manufacturer's protocol. cDNA (cDNA) was synthesized from 1 µg of total RNA using Superscript III Reverse Transcriptase (Invitrogen). PCR was carried out using ABI StepOne real-time PCR system as the following steps: 95°C for 5 min, followed by 50 cycles of successive incubation at 95°C for 30 sec, 62°C for 15 sec, and 68°C for 30 sec. *FLT4* cDNA was amplified with the following primers: forward, 5'-TGTGGAGGGAAAGAATAAGACTGT-3'; reverse, 5'-GCCCCACCTTGTGGAGAC-3'. *GAPDH* cDNA was amplified with the following primers: forward, 5'-

TTGACGGTGCCATGGAATTT-3'; reverse, 5'-GCCATCAATGACCCCTTCATT-3'. The expression of *FLT4* was normalized against that of *GAPDH*.

#### Statistical analyses

Statistical analyses were performed using SAS v9.3 (SAS Institutes), and the heat maps were generated by the gplots package in R.

#### Selection of biomarkers for detecting tumor and normal samples

The AUC was calculated for each CpG site to evaluate the accuracy. Leave-one-out cross validation was conducted to test the reliability without collecting new samples. Moreover, a subset analysis was conducted for the ability to distinguish early-stage OSCC (n=16) and normal (n=15) samples.

#### Cluster analysis of methylation levels

The normalized methylation levels in  $\beta$  values for profiled CpG sites were presented in a heat map and analyzed with a hierarchical agglomerative cluster method for differential methylation between tumor and normal samples. In the cluster analysis, we used Manhattan distance to measure the distance between observed values and defined clusters by the method of complete linkage.

#### Validation of *FLT4* methylation status and RNA expression

Quantitative methylation levels between different types of tumor and normal samples measured by pyrosequencing assay were compared by Mann-Whitney U test. Wilcoxon signed rank test was used to test the difference in fold-change of RNA expression.

#### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

#### Funding

This study was supported in part by the Ministry of Science and Technology of Taiwan (MOST103-2314-B-039-037 and NSC99-2320-B-194-004-MY3).

#### References

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; 136:E359-86; PMID:25220842; <http://dx.doi.org/10.1002/ijc.29210>
2. Department of Health, Executive Yuan, R.O.C.(TAIWAN). 2009 Statistics of Causes of Death. Taipei, Taiwan, 2010.
3. Warnakulasuriya S. Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol* 2009; 45:309-16; PMID:18804401; <http://dx.doi.org/10.1016/j.oraloncology.2008.06.002>
4. Lo WL, Kao SY, Chi LY, Wong YK, Chang RC. Outcomes of oral squamous cell carcinoma in Taiwan after surgical therapy: factors affecting survival. *Journal of Oral and Maxillofacial Surgery* 2003; 61:751-8; PMID:12856245; [http://dx.doi.org/10.1016/S0278-2391\(03\)00149-6](http://dx.doi.org/10.1016/S0278-2391(03)00149-6)
5. Richards D. Patient delay in reporting oral cancer is poorly understood. *Evid Based Dent* 2007; 8:21; PMID:17380180; <http://dx.doi.org/10.1038/sj.ebd.6400472>
6. Chen CL, Chi CW, Chang KW, Liu TY. Safrole-like DNA adducts in oral tissue from oral cancer patients with a betel quid chewing history. *Carcinogenesis* 1999; 20:2331-4; PMID:10590228; <http://dx.doi.org/10.1093/carcin/20.12.2331>
7. Ko YC, Huang YL, Lee CH, Chen MJ, Lin LM, Tsai CC. Betel quid chewing, cigarette smoking and alcohol consumption related to oral cancer in Taiwan. *J Oral Pathol Med* 1995; 24:450-3; PMID:8600280; <http://dx.doi.org/10.1111/j.1600-0714.1995.tb01132.x>
8. Jones PA. DNA methylation and cancer. *Oncogene* 2002; 21:5358-60; PMID:12154398; <http://dx.doi.org/10.1038/sj.onc.1205597>
9. Shaw RJ, Akufo-Tetteh EK, Risk JM, Field JK, Lilo-glou T. Methylation enrichment pyrosequencing: combining the specificity of MSP with validation by pyrosequencing. *Nucleic Acids Res* 2006; 34:e78; PMID:16807314; <http://dx.doi.org/10.1093/nar/gkl424>
10. Viet CT, Schmidt BL. Methylation array analysis of preoperative and postoperative saliva DNA in oral cancer patients. *Cancer Epidemiol Biomarkers Prev* 2008; 17:3603-11; PMID:19064577; <http://dx.doi.org/10.1158/1055-9965.EPI-08-0507>



11. Viswanathan M, Tsuchida N, Shanmugam G. Promoter hypermethylation profile of tumor-associated genes p16, p15, hMLH1, MGMT and E-cadherin in oral squamous cell carcinoma. *Int J Cancer* 2003; 105:41-6; PMID:12672028; <http://dx.doi.org/10.1002/ijc.11028>
12. Carvalho AL, Jeronimo C, Kim MM, Henrique R, Zhang Z, Hoque MO, Chang S, Brait M, Nayak CS, Jiang WW, et al. Evaluation of promoter hypermethylation detection in body fluids as a Screening/Diagnosis tool for head and neck squamous cell carcinoma. *Clinical Cancer Research* 2008; 14:97-107; PMID:18172258; <http://dx.doi.org/10.1158/1078-0432.CCR-07-0722>
13. Mosch B, Reissenweber B, Neuber C, Pietzsch J. Eph receptors and ephrin ligands: important players in angiogenesis and tumor angiogenesis. *J Oncol* 2010; 2010:135285; PMID:20224755
14. Hu L, Sham JS, Xie D, Wen JM, Wang WS, Wang Y, Guan XY. Up-regulation of fibroblast growth factor 3 is associated with tumor metastasis and recurrence in human hepatocellular carcinoma. *Cancer Lett* 2007; 252:36-42; PMID:17215076; <http://dx.doi.org/10.1016/j.canlet.2006.12.003>
15. Ruscica M, Dozio E, Motta M, Magni P. Relevance of the neuropeptide Y system in the biology of cancer progression. *Curr Top Med Chem* 2007; 7:1682-91; PMID:17979777; <http://dx.doi.org/10.2174/156802607782341019>
16. Kitlinska J. Neuropeptide Y in neural crest-derived tumors: effect on growth and vascularization. *Cancer Lett* 2007; 245:293-302; PMID:16513255; <http://dx.doi.org/10.1016/j.canlet.2006.01.010>
17. Ju YT, Chang AC, She BR, Tsaur ML, Hwang HM, Chao CC, Cohen SN, Lin-Chao S. gas7: A gene expressed preferentially in growth-arrested fibroblasts and terminally differentiated Purkinje neurons affects neurite formation. *Proc Natl Acad Sci U S A* 1998; 95:11423-8; PMID:9736752; <http://dx.doi.org/10.1073/pnas.95.19.11423>
18. Ball DW. Achaete-scute homolog-1 and Notch in lung neuroendocrine development and cancer. *Cancer Lett* 2004; 204:159-69; PMID:15013215; [http://dx.doi.org/10.1016/S0304-3835\(03\)00452-X](http://dx.doi.org/10.1016/S0304-3835(03)00452-X)
19. Chang JT, Lu YC, Chen YJ, Tseng CP, Chen YL, Fang CW, Cheng AJ. hTERT phosphorylation by PKC is essential for telomerase holoprotein integrity and enzyme activity in head neck cancer cells. *Br J Cancer* 2006; 94:870-8; PMID:16508638; <http://dx.doi.org/10.1038/sj.bjc.6603008>
20. Sherwood NM, Krueckl SL, McRory JE. The origin and function of the pituitary adenylate cyclase-activating polypeptide (PACAP)/glucagon superfamily. *Endocr Rev* 2000; 21:619-70; PMID:11133067
21. Hayes JD, McLellan LI. Glutathione and glutathione-dependent enzymes represent a co-ordinately regulated defence against oxidative stress. *Free Radic Res* 1999; 31:273-300; PMID:10517533; <http://dx.doi.org/10.1080/10715769900300851>
22. Thirumoorthy N, Manisenthil Kumar KT, Shyam Sundar A, Panayappan L, Chatterjee M. Metallothionein: an overview. *World J Gastroenterol* 2007; 13:993-6; PMID:17373731; <http://dx.doi.org/10.3748/wjg.v13.i7.993>
23. Chand HS, Schmidt AE, Bajaj SP, Kiesel W. Structure-function analysis of the reactive site in the first Kunitz-type domain of human tissue factor pathway inhibitor-2. *J Biol Chem* 2004; 279:17500-7; PMID:14970225; <http://dx.doi.org/10.1074/jbc.M400802200>
24. Petersen LC, Sprecher CA, Foster DC, Blumberg H, Hamamoto T, Kiesel W. Inhibitory properties of a novel human Kunitz-type protease inhibitor homologous to tissue factor pathway inhibitor. *Biochemistry* 1996; 35:266-72; PMID:8555184; <http://dx.doi.org/10.1021/bi951501d>
25. Konduri SD, Tasiou A, Chandrasekar N, Rao JS. Overexpression of tissue factor pathway inhibitor-2 (TFPI-2), decreases the invasiveness of prostate cancer cells in vitro. *Int J Oncol* 2001; 18:127-31; PMID:11115549
26. Lai YH, He RY, Chou JL, Chan MW, Li YF, Tai CK. Promoter hypermethylation and silencing of tissue factor pathway inhibitor-2 in oral squamous cell carcinoma. *Journal of translational medicine* 2014; 12:237; PMID:25179542; <http://dx.doi.org/10.1186/s12967-014-0237-7>
27. Wong CM, Ng YL, Lee JM, Wong CC, Cheung OF, Chan CY, Tung EK, Ching YP, Ng IO. Tissue factor pathway inhibitor-2 as a frequently silenced tumor suppressor gene in hepatocellular carcinoma. *Hepatology* 2007; 45:1129-38; PMID:17464989; <http://dx.doi.org/10.1002/hep.21578>
28. Ferrara N, Davis-Smyth T. The biology of vascular endothelial growth factor. *Endocr Rev* 1997; 18:4-25; PMID:9034784; <http://dx.doi.org/10.1210/edrv.18.1.0287>
29. Ferrara N, Bunting S. Vascular endothelial growth factor, a specific regulator of angiogenesis. *Curr Opin Nephrol Hypertens* 1996; 5:35-44; PMID:8834160; <http://dx.doi.org/10.1097/00041552-199601000-00008>
30. Klagsbrun M, D'Amore PA. Vascular endothelial growth factor and its receptors. *Cytokine Growth Factor Rev* 1996; 7:259-70; PMID:8971481; [http://dx.doi.org/10.1016/S1359-6101\(96\)00027-5](http://dx.doi.org/10.1016/S1359-6101(96)00027-5)
31. Veikkola T, Alitalo K. VEGFs, receptors and angiogenesis. *Semin Cancer Biol* 1999; 9:211-20; PMID:10343072; <http://dx.doi.org/10.1006/scbi.1998.0091>
32. Guan M, Xu C, Zhang F, Ye C. Aberrant methylation of EphA7 in human prostate cancer and its relation to clinicopathologic features. *Int J Cancer* 2009; 124:88-94; PMID:18821581; <http://dx.doi.org/10.1002/ijc.23890>
33. Wang J, Kataoka H, Suzuki M, Sato N, Nakamura R, Tao H, Maruyama K, Isogaki J, Kanaoka S, Ihara M, et al. Downregulation of EphA7 by hypermethylation in colorectal cancer. *Oncogene* 2005; 24:5637-47; PMID:16007213; <http://dx.doi.org/10.1038/sj.onc.1208720>
34. Peng DF, Razvi M, Chen H, Washington K, Roessner A, Schneider-Stock R, El-Rifai W. DNA hypermethylation regulates the expression of members of the Mu-class glutathione S-transferases and glutathione peroxidases in Barrett's adenocarcinoma. *Gut* 2009; 58:5-15; PMID:18664505; <http://dx.doi.org/10.1136/gut.2007.146290>
35. Yu J, Zhu T, Wang Z, Zhang H, Qian Z, Xu H, Gao B, Wang W, Gu L, Meng J, et al. A novel set of DNA methylation markers in urine sediments for sensitive/specific detection of bladder cancer. *Clin Cancer Res* 2007; 13:7296-304; PMID:18094410; <http://dx.doi.org/10.1158/1078-0432.CCR-07-0861>
36. Bibikova M, Lin Z, Zhou L, Chudin E, Garcia EW, Wu B, Doucet D, Thomas NJ, Wang Y, Vollmer E, et al. High-throughput DNA methylation profiling using universal bead arrays. *Genome Res* 2006; 16:383-93; PMID:16449502; <http://dx.doi.org/10.1101/gr.4410706>
37. Gigeck CO, Leal MF, Silva PN, Lisboa LC, Lima EM, Calcagno DQ, Assumpcao PP, Burbano RR, Smith Mde A. hTERT methylation and expression in gastric cancer. *Biomarkers* 2009; 14:630-6; PMID:20001710; <http://dx.doi.org/10.3109/13547500903225912>
38. Clement G, Braunschweig R, Pasquier N, Bosman FT, Benhattar J. Methylation of APC, TIMP3, and TERT: a new predictive marker to distinguish Barrett's oesophagus patients at risk for malignant transformation. *J Pathol* 2006; 208:100-7; PMID:16278815; <http://dx.doi.org/10.1002/path.1884>
39. Jithesh PV, Risk JM, Schache AG, Dhanda J, Lane B, Liloglou T, Shaw RJ. The epigenetic landscape of oral squamous cell carcinoma. *Br J Cancer* 2013; 108:370-9; PMID:23287992; <http://dx.doi.org/10.1038/bjc.2012.568>
40. Nagata S, Hamada T, Yamada N, Yokoyama S, Kitamoto S, Kanmura Y, Nomura M, Kamikawa Y, Yonezawa S, Sugihara K. Aberrant DNA methylation of tumor-related genes in oral rinse: a noninvasive method for detection of oral squamous cell carcinoma. *Cancer* 2012; 118:4298-308; PMID:22252571; <http://dx.doi.org/10.1002/cncr.27417>
41. Richards KL, Zhang B, Baggerly KA, Colella S, Lang JC, Schuller DE, Krahe R. Genome-wide hypomethylation in head and neck cancer is more pronounced in HPV-negative tumors and is associated with genomic instability. *PLoS One* 2009; 4:e4941; PMID:19293934; <http://dx.doi.org/10.1371/journal.pone.0004941>
42. Sartor MA, Dolinoy DC, Jones TR, Colacino JA, Prince ME, Carey TE, Rozek LS. Genome-wide methylation and expression differences in HPV(+) and HPV(-) squamous cell carcinoma cell lines are consistent with divergent mechanisms of carcinogenesis. *Epigenetics* 2011; 6:777-87; PMID:21613826; <http://dx.doi.org/10.4161/epi.6.6.16216>
43. Shaw R, Robinson M. The increasing clinical relevance of human papillomavirus type 16 (HPV-16) infection in oropharyngeal cancer. *Br J Oral Maxillofac Surg* 2011; 49:423-9; PMID:20727631; <http://dx.doi.org/10.1016/j.bjoms.2010.06.023>
44. Chang KC, Su IJ, Tsai ST, Shieh DB, Jin YT. Pathological features of betel quid-related oral epithelial lesions in taiwan with special emphasis on the tumor progression and human papillomavirus association. *Oncology* 2002; 63:362-9; PMID:12417791; <http://dx.doi.org/10.1159/000066227>
45. Righini CA, de Fraipont F, Timsit JF, Faure C, Brambilla E, Rey E, Favrot MC. Tumor-specific methylation in saliva: a promising biomarker for early detection of head and neck cancer recurrence. *Clin Cancer Res* 2007; 13:1179-85; PMID:17317827; <http://dx.doi.org/10.1158/1078-0432.CCR-06-2027>